

INDEX

Note: AF stands for atrial fibrillation.

Page numbers in *italics* indicate figures.

Page numbers followed by a “t” indicate tables.

Clinical trials and studies are indexed
under the acronym of the name.

A to Z Trial, 95

AADs. See *Antiarrhythmic drugs (AADs)*.

Ablation. See also *Catheter ablation; Surgical ablation*.

catheter ablation, 13, 164-184

success rate, 133

clinical studies, 118t-119t

EPS for targeting sites for, 79

with pacemaker implantation, 270-272

surgical ablation, 184-187, 185, 186t, 270-272, 279, 381

ACC/AHA/ESC Guidelines

AF classification, 17-20

diagnostic tests, 72t-73t

pharmacologic cardioversion, 152

ACC/AHA/HRS Guidelines

ADDS for repeat cardioversion, 145, 145t

antithrombotic therapy, 289, 295

catheter ablation, 181-184

device guidelines, 103-104

electrical cardioversion, 134-135

prevention of AF, 128

ACE inhibitors, 88-93, 89, 104, 305

actions of, 90t

in combination with AADs, 214

Acetabutolol, 260t

Acetaminophen, 317, 318t

Acetylcholine, 40

ACT I-IV trials, 161-163

ACTIVE-A study, 313, 314t, 358

ACTIVE-W trial, 310, 358

Acute decompensated heart failure (ADHF), 65

ACUTE trial, 361-362

ACUTE II study, 363

Adenosine, 57

ADONIS, 226, 229

Adrenergic tone, high, 258

AF-CHF trial, 122, 126-127

AF Symptoms Study, 272

AFASAK, 293, 304

AFFECTS Registry, 311

AFFIRM trial, 31, 116-123, 118t-119t, 120, 126, 128, 194, 216, 221, 254-255, 265, 270
 hazard ratios for death in subgroups, 121
 AFI trial, 294
 African Americans, 27, 52, 85
 Age. See also *Elderly patients; Young patients.*
 AF prevalence and incidence and, 23-25, 24, 26, 27, 61
 AF risk and, 49, 61
 in CHA2DS2-VASc score, 298t
 in CHADS2 score, 294, 294t
 choice of therapy and, 124, 127, 128, 129t, 133
 degeneration of conduction system and, 270
 hospitalizations for AF, 30
 thromboembolic risk and, 286, 292, 293t
 Aggrastat to Zocor (A to Z) Trial, 95
 AHA/ACC/HRS guidelines (2014), 289
 Albuterol, 57
 Alcohol intake, 56, 73, 320-321
 Aldosterone, 44
 Aldosterone antagonists, 88, 90t, 93
 Allopurinol, 319t
 Ambulatory arrhythmia monitoring, 77, 79-80, 81t
 American College of Cardiology. See *ACC/AHA/ESC Guidelines*.
 American Heart Association. See *ACC/AHA/ESC Guidelines*.
 Amiodarone, 116, 159t, 207t, 216-222
 action mechanisms, 207t, 216, 266
 adverse effects, 207t, 209, 218-219, 267
 clinical studies, 118t-119t, 125, 126, 220-222
 contraindications/cautions, 219-220, 324
 derivative of. See *Dronedarone*.
 dosing, 159t, 207t, 209, 217-218
 low-dose, 212
 drug/device interactions, 159t, 219-220, 264, 267, 318t
 implanted device threshold effects, 199t
 indications for, 159t, 216-222
 cardioversion
 failed electrical, 147, 148t
 pharmacologic, 151, 159t
 pretreatment for, 148t, 149
 repeat electrical, 143, 143t, 146t
 maintenance of NSR, 207t, 216-222
 rate control, 258, 266-267
 initiation of therapy, 213, 215t
 monitoring, 201, 202t-203t, 219
 pharmacokinetics, 216
 safety of, 159t, 198, 199, 201-204, 207t, 209, 212, 218-222
 hypotension, 147
 organ toxicity, 209, 212
 toxicity with long-term use, 127, 200, 209, 212

Amoxicillin, 318t
 Amplatzer Cardiac Plug, 372, 373, 379-381
 Anabolic steroids, 319t
 ANDROMEDA, 227
 Anesthesia, for electrical cardioversion, 135-136
 Angiotensin-converting enzyme inhibitors. See *ACE inhibitors*.
 Angiotensin II, 44
 Angiotensin receptor blockers (ARBs), 88-93, 89, 104
 actions of, 90t
 ANP (atrial natriuretic peptide), 43, 60
 Anti-inflammatory agents, 40, 93, 98, 317, 318t
 Antiarrhythmic drugs (AADs), 193-252
 action mechanisms, 204, 206t-207t
 adverse events, 206t-207t
 agents available, 156t-159t, 206t-207t, 216-248
 amiodarone, 207t, 216-222
 disopyramide, 206t, 246-248
 dofetilide, 207t, 237-241
 dronedarone, 207t, 222-233
 ideal agent, 155-164, 205
 propafenone, 206t, 242-246
 sotalol, 207t, 233-236
 anticoagulation and, 193, 213, 215-216
 atrial selective, 155-164
 benefits of, questions on, 115
 cardioversion and, 150-164
 electrical cardioversion
 failed, 147-148, 148t
 repeat, 145, 145t
 pharmacologic cardioversion, 150-164
 choice of AADs for, 154t
 new (atrial selective) AADs, 155-164
 pretreatment prior to cardioversion, 145, 145t, 148-149
 choice of, 204-212, 206t-207t, 210-211
 algorithm/stepwise approach, 208
 atrial selective agents, 155-164
 available agents, 216-248
 guidelines and recommendations, 205, 208, 210-211
 heart failure patients, 212
 hypertensive patients, 209
 patients with coronary artery disease, 209-210
 patients with no or minimal heart disease, 205-209
 for pharmacologic cardioversion, 152, 153t, 154t, 156t-159t
 classification (Vaughan-Williams), 206t-207t
 clinical trials, 116-124, 118t-119t, 220-248
 combination therapy, 214-215
 dosing, 206t-207t
 booster dosing, 213
 drug interactions, 318t

Antiarrhythmic drugs (AADs) (*continued*)

efficacy, 115-116, 118t-119t, 119t, 194
goals of therapy, 193
indications for, 194
 for rate control, 258, 264, 266-267
initiation of therapy, 205, 212-214, 278-279
 hospitalization requirements for, 213, 215t
test prior to starting, 77-78
for maintenance of NSR, 133, 183-252
agents available, 206t-207t
choice of AAD, 204-223, 206t-207t, 208, 210-211
efficacy, 194
general principles, 193-200
goals of therapy, 193
indications for, 194
initiation of therapy, 205, 212-214
safety considerations, 193, 195-198
stepwise approach, 208
monitoring, 80, 200-204, 202t-203t
 of initiation phase, 212-214, 215t
outpatient use of, 213, 215t
for pharmacologic cardioversion, 154t
pill-in-the-pocket approach, 151, 213
pretreatment with, prior to cardioversion, 145, 145t, 148-149
for rate control, 258, 264, 266-267
safety considerations, 117-120, 151-152, 194, 195-198
 AF conversion to atrial flutter, 152
 atrial proarrhythmias, 197-198
 atrial remodeling and, 165
blood tests/monitoring, 79, 80, 200-204, 202t-203t
bradyarrhythmia, 197, 198, 212
hemodynamic effects, 199-200
implanted device function, 199, 199t
mortality, 117-120
organ toxicity, 200, 209, 212
proarrhythmias, 195-198, 209
QT prolongation, 196-197, 198t
risk-benefit ratio, 205
torsade de pointes (TdP), 195, 196-197, 196, 213
toxicity, 115, 127, 129, 151-152, 200, 209
 ventricular proarrhythmias, 195-197, 196, 212
side effects, 195
vs catheter ablation, 175-177, 178t-180t, 182-183
Antibiotics, 198t, 317, 318t
Anticoagulation therapy, 151, 193, 304-372. See also *Warfarin*.
agents available
 AADs, 193, 213, 215-216
 apixaban, 339-347
 dabigatran etexilate, 323-334

Anticoagulation therapy, agents available (*continued*)

direct thrombin inhibitors, 304, 305, 322-334
 edoxaban, 334, 347-351
 factor Xa inhibitors, 304, 305, 334-351
 rivaroxaban, 335-339
 warfarin, 304-322, 356t
bridging therapy, 369, 370
with cardioversion procedures, 136-137, 151, 359-363, 366-367
 AF duration and, 359-363
 prevention of new thrombus, 363, 364
 risk of preexisting thrombus, 360-363
for catheter ablation, 170-171
with catheter ablation, 171
 chronic, 289
 discontinuance prior to catheter ablation, 171
duration of treatment with, 121, 124, 129, 134
long-term continuation of, 121, 129
newer anticoagulants, 304, 322-357
 advantages and disadvantages, 356t
 direct thrombin inhibitors, 304, 305, 322-334
 factor Xa inhibitors, 304, 334-351
 summary of, 351-357, 356t
oral anticoagulants, 304-322
prophylactic, for cardioversion, 136-137
special situations
 AF ablation and Cox-Maze procedure, 364
 AF due to reversible causes, 368-369
 AF patients presenting with stroke, 364-365
 atrial flutter, 365-368
 cardioversion procedures, 359-363
 interruption of anticoagulation in AF patients, 369
 patients with coronary interventions, 369-372
 warfarin interruption, 370
for stroke prevention, 170-171
 efficacy, 116, 120-121
 risks vs benefits, 289
 treatment algorithm, 305
in treatment algorithm, 305
triple therapy, 369-370, 371-372
Antidepressants, 198t
Antifungal agents, 198t
Antihistamines, 198t, 371
Antihypertensive agents, 61, 305
Antioxidants, 101
Antiplatelet therapy, 113, 305, 357-359
 aspirin, 357-358
bleeding risk, 358, 370-372
combination of agents, 304, 358-359
dosage, 371

Antiplatelet therapy (*continued*)
 drug interactions, 318t
 efficacy of, 304
 Antithrombotic therapy, 113, 296t, 303-372.
 See also *Thromboembolism*.
 ACCP guidelines (2014), 295, 296t
 Antithyroid drugs, 319t
 Antiviral agents, 198t, 318t
 Anxiety, differential diagnosis of, 67t
 Aortic plaque, 292
 Aortic stenosis, 65, 68
 Apixaban, 335, 339-347
 as alternative to warfarin, 295, 351
 bleeding risk, 331-334, 333t
 clinical trials, 339-347, 343t, 344
 ARISTOTLE, 342-347, 346t, 347, 348t
 AVEROES, 339-342, 343t, 344
 in treatment algorithm, 305
 Aprindine, 126
 ARBs (angiotensin receptor blockers), 88-93, 89, 104, 305
 in combination with AADs, 214
 ARISTOTLE study, 342-347, 346t, 347, 348t
 ARMYDA3 trial, 94
 Arrhythmias
 related, 16-17, 17, 48t
 shock-induced, 134, 149
 ASA. See *Aspirin*.
 ASAP study, 379
 Aspirin, 357-358
 drug interactions, 317
 SPAF trials, 357-358
 stroke hazard rate with, 357
 for stroke prevention, 296t, 357-358
 combination therapy, 304, 305, 358-359
 in treatment algorithm, 305
 in triple therapy, 369-370, 371-372
 Astemizole, 198t
 Asymptomatic AF, 31, 65, 69
 treatment choice and, 129, 129t
 Atenolol, 258, 260t, 265, 266
 ATHENA trial, 227-229, 230-231
 Atherosclerotic plaque, 283
 Athletes, 51, 52
 Atorvastatin, 94-98, 96t-97t
 Atria, electrophysiologic properties of, 155, 160
 ATRIA study, 23, 314, 315
 Atrial arrhythmias, 48t
 Atrial defibrillators, 134, 135t, 140, 149-150

Atrial dilatation, 47, 48t, 53, 284
 Atrial electrical abnormalities, 34
 Atrial-esophageal fistula, 172, 173t
 Atrial fibrillation (AF)
 classification of, 17-20
 consequences and prognosis of, 27-31
 definition of, 13
 on electrocardiogram (ECG), 15
 epidemiology, 23-32
 mechanism of, 33-40, 34
 monitoring of, 76-80, 81t
 new-onset, stroke risk in, 300-301, 302
 pathophysiology of, 33-45, 34
 perpetuation of (“AF begets AF”), 38-39, 85, 193
 prevalence and incidence of, 23-26, 24, 25
 prevention of, 83-108, 84
 related arrhythmias, 16-17, 17
 therapeutic approaches, 112-113. See also *Therapy*.
 triggers of, 17, 33-36, 167
 variety of clinical presentations in, 65, 69
 Atrial fibrosis. See *Fibrosis*.
 Atrial flutter, 16-17, 17, 57-58
 after catheter ablation, 174
 anticoagulation for, 365-368
 cardioversion of, 164
 conversion of AF to, 139, 152
 electrocardiogram of, 18
 EPS for, 79
 Atrial natriuretic peptide (ANP), 43, 60
 Atrial pressure, elevated, 47, 48t
 Atrial remodeling, 39, 85-88, 150, 165
 anti-remodeling therapies, 85-88, 89
 regression with AADs, 193
 Atrial selective AADs, 155-160
 Atrial septal defect (ASD), 54
 Atrial size/volume, 36, 60, 75
 Atrial structural abnormalities, 34
 Atrial stunning, 136, 142-143, 284
 Atrial systolic function, 142-143
 Atrial tachycardia, 16, 17, 17. See also *Tachycardia*.
 Autoimmune processes in AF, 38
 Autonomic nervous system/tone, 34, 40, 47, 48t, 204
 AV junction RV ablation, 118t-119t
 AV nodal ablation, 270-272, 277, 279
 risks and complications of, 271-272
 AV nodal agents, 274. See also *Rate control*.
 AV nodal block, 277
 AV nodal modification, 272

- AV nodal vagal stimulation, 273
 AVERROES trial, 339-342, 343t, 344
 Azathioprine, 318t
- BAATAF, 293, 304
 BAFTA study, 309
 Barbiturates, 319t
 Baylor University Medical Center study, 58-59, 59
 β-Agonists, 54
 β-Blockers, 74, 89, 258-260
 clinical studies, 117, 118t-119t, 126
 combination therapy, 274, 275, 278
 with statins, 95-98
 contraindications/cautions, 258-260, 275
 indications for, 56, 128, 152, 205, 209, 258-260
 for rate control, 255, 258-260, 259t, 260t, 273-274, 278
 Birmingham scheme of stroke risk, 298, 298t, 300, 309
 Bleeding risk, 289, 306
 antiplatelet therapy, 358
 apixaban, 331-334, 333t
 dabigatran, 337-339, 340t-341t
 edoxaban, 350-351, 354t-355t
 strategies to reduce, 370-372
 vs stroke prevention, 315
 warfarin, 306, 313-315, 316, 317, 326-329, 328t
- Blood pressure, 61
 hypertension, 51-52. See also *Hypertension*.
 hypotension, 65
- Blood stasis, thromboembolism and, 284
- Blood tests, 71, 72t, 76-77, 80
- Blue toe syndrome, 321
- BNP. See *Brain natriuretic peptide*.
- Body mass index (BMI), 55, 61
- Bosentan, 318t
- Bradyarrhythmia, 149, 198
- Bradycardia, 142, 152, 197, 198, 212
 rate-control therapy and, 256-257
- Brain imaging, 287
- Brain natriuretic peptide (BNP), 43-44, 60, 79
- Bypass surgery. See *CABG*.
- C-reactive protein (CRP), 38, 60, 95, 98, 285
- CABANA trial, 177
- CABG, post-surgery AF and, 48t, 55, 57, 58-59, 59
 anticoagulation, 368-369, 370
 case study, 393-395
 electrical cardioversion and, 141
 steroid therapy and, 98-101, 100
- CAFA, 293, 304, 306
- Caffeine, 26, 56-57
 Calcium channel blockers (CCBs), 89, 260-261
 combination therapy, 265, 266, 274, 275, 278
 contraindications and cautions, 260, 275
 for rate control, 128, 152, 255, 260-261, 265, 273-274, 275, 278
 for repeat cardioversion, 145
 Calcium loading, 89
 Canadian Cardiovascular Society Severity in Atrial Fibrillation (CCS-SAF) scoring system, 73
 Canadian trials. See *CTAF; CTOPP*.
- Candesartan, 90-91
- CAPP, 91
- CARAF I and II studies, 95-98, 245-246
- Carbamazepine, 319t
- Cardiac Arrhythmia Suppression Trial, 245
- Cardiac conditions associated with AF, 48t, 51-54
- Cardiac murmur, 61
- Cardiac pacemakers. See *Pacemakers*.
- Cardiac surgery, post-operative AF and, 40, 57, 58-59, 59
 electrical cardioversion and, 142
 steroid therapy and, 98-101, 100
- Cardiac tamponade, 172-173, 173t
- Cardiac thromboembolism, 285
- Cardioembolic stroke, 286-287, 288
- Cardiomyopathy, tachycardia-induced, 28, 41, 42, 65, 69, 128
- Cardiovascular disease, 25
- Cardiovascular drugs, 318t
- Cardiovascular Health Study (CHS), 23
- Cardiovascular risk factors for thromboembolism, 293t
- Cardiovascular side effects of AAD initiation, 212-213
- Cardioversion, 115, 133-192. See also *Electrical cardioversion; Pharmacologic cardioversion*.
 anticoagulation for, 136-137, 151, 359-363, 366-367
 prevention of new thrombus, 363, 364
 risk of preexisting thrombus, 360-363
 atrial defibrillators, 134, 135t, 140, 149-150
 checklist for prior to, 134, 135t
 direct current, 134
 electrical, 134-151
 safety of, 151
 failed, 143-144, 144, 146
 pharmacologic facilitation of, 147-149, 148t
 high energy shock, 139, 147t
 indications for, 134-135, 137, 208, 275, 277
 internal, 133, 134, 149
 overview of, 134
 pharmacologic, 150-164
 factors to consider, 154, 154t
 new AADs for, 155-164, 160

- Cardioversion, pharmacologic (*continued*)
 pill-in-the-pocket strategy, 151, 158t, 213
 recommended AADs for, 154t, 156t-159t
 pretreatment with AADs prior to, 145, 145t, 148-149
 repeat, AADs for, 145, 145t, 146t, 147
 spontaneous, 150
 TEE-guided, 80, 361-363
 thromboembolism after, 285, 359-363
CARTO, 170
Carvedilol, 260t
Case studies, 389-399
 AF patient with stent placement, 397-399
 AF post-CABG, 393-395
 AF with hyperthyroidism, 395-397
 elderly patient with new-onset AF, 391-393
 lone AF, 389-391
CAST, 77, 115, 212, 245
Catecholamines, 35
 Catheter ablation, 133, 164-184
 adverse events, 176
 clinical trials, 178t-180t
 complications, 171-173
 efficacy and safety, 151, 176-177, 181
 focal triggers, 165, 166
 follow-up and recurrence of AF, 173-174
 indications for, 133-134, 151, 177-184
 outcomes and further management, 175
 periprocedural management, 170-171
 anticoagulation, 170-171
 imaging, 170
 TEE, 170, 171
 repeat procedure, 174
 safety and efficacy, 176-177, 178t-180t, 181-184, 181
 success rate, 133
 techniques and procedures, 166-170, 168
 vs AADs, 175-177, 178t-180t, 182-183
 vs antiarrhythmic therapy, 175-177
CCBs. See *Calcium channel blockers*.
Cephalosporins, 318t
Cerebral thromboembolism, 286
Cerebrovascular accident (CVA), 65, 69, 172
Cesarean section, 322
CHA2DS2-VASc, 295, 297, 298-300, 298t
 compared with CHADS2 score, 299, 299t
CHADS2 score, 60, 171, 294-297, 294t, 300
 compared with CHA2DS2-VASc score, 299, 299t
 limitations of, 295
 predictive value of, 295
CHARM trial, 90-91
CHAS2DS2-VASc score, 298t
Chest imaging, 71, 73t, 77
Chest pain, 43, 66t-67t
Chloroquine, 198t
Cholesterol embolization, 321
Cholestyramine, 318t
Chronic medical conditions, 48t, 54-57
Chronic obstructive pulmonary disease (COPD), 54
Chronotropic incompetence, 257
Cibenzoline, 126
Cigarette smoking, 57, 83
Cimetidine, 238, 240, 319t
Ciprofloxacin, 317
Circumferential left atrial ablation, 168, 169-170
Cisapride, 198t
Clarithromycin, 317, 335
Classification of AF, 17-20, 68
 episodes shorter than 30 seconds, 20
 first-detected episode, 19
 lone AF, 20, 47
 long-standing persistent AF, 19
 non-valvular AF, 19
 paroxysmal AF, 19
 permanent AF, 19
 persistent AF, 19
 recurrent episode, 19
 secondary to other conditions, 20
Clinical manifestations, 65-70
 asymptomatic AF, 31, 65, 69
 differential diagnosis of chest pain, 66t-67t
 poor correlation with AF, 68, 78
 variation in, 65, 69
Clonidine, 267-268, 278
Clopidogrel, 305, 358-359
 in combination with aspirin, 305, 358-359, 370-371
 in combination with irbesartan, 310
 drug interactions, 317
CNs agents, 319t
Coagulation. See also *Anticoagulation therapy*.
 anticoagulation therapy, 151, 193, 304-322
 hypercoagulable state, 28, 284-285
Coffee consumption, 26, 56-57
Complications
 of AF, 65, 69, 84
 of catheter ablation, 171-173
 of DC cardioversion, 142-143
 thromboembolic, 120-121, 143, 285, 288, 290t-291t
Computed tomography. See *CT scans*.
Conduction block, 35

Congenital heart disease, 54
 Consequences of AF, 27-31
 hospitalizations and costs, 29-31, 30
 impaired hemodynamics, 28-29
 mortality, 27, 29
 quality of life, reduced, 27, 31
 thromboembolism, 27-28
 COPD (chronic obstructive pulmonary disease), 54
 Copenhagen Stroke Study, 28
 COPPA II, 246
 Coronary artery bypass graft. See *CABG*.
 Coronary artery disease (CAD), 47, 53, 66t
 AADs for, 209-212
 Coronary blood flow, impaired, 43
 Coronary interventions, anticoagulation for, 369-372
 Coronary stents, 371-372
 Corticosteroids. See *Glucocorticosteroids*.
 Cost-effectiveness of treatment, 124
 Costs of AF, 29-31
 Cox, James, 184
 Cox-Maze procedure, 184-187, 185. See also *Surgical ablation*.
 COX-2 inhibitors, 317, 318t
 CRAFT, 161
 Creatinine, serum, 223
 CRRAFT, 127
 CT scans, 77, 170
 CTAF, 122, 221, 222, 235-236, 245
 CTOPP, 103
 CYP systems, 161, 316

Dabigatran etexilate, 323-334
 as alternative to warfarin, 295, 351
 bleeding and, 337-339, 340t-341t
 characteristics, 323-325
 clinical trials, 325-331
 contraindications and cautions, 324
 dosing/interactions, 324
 efficacy and safety, 324-331
 FDA approval of, 323
 RE-LY trial, 325-330, 326, 327t
 RELY-ABLE trial, 330-331
 risk category and, 296t
 switching to and from, 324-325
 in treatment algorithm, 305

DAFNE, 225-226

Danish Diet, Cancer, and Health Study, 57, 102

DC cardioversion, 134

Decongestants, 74

Defibrillation, atrial, 134, 135t, 140, 149-150

Defibrillators, 141, 149-150
 dual-chamber, 149-150
 memory function of, 81t
 Definition of AF, 13
 Demographics of AF, 27
 Device-based therapy, 103-104, 113. See also *Pacemakers*.
 Device interactions. See *specific agents*.
 Dexamethasone, 99
 Diabetes, 20
 in CHA2DS2-VASc score, 298t
 in CHADS2 score, 294, 294t
 Diagnosis of AF. See also *Classification of AF; Evaluation of patient with AF*.
 characterization of episodes, 17-19
 clinical manifestations, 65-70
 differential diagnosis, 16, 66t-67t
 electrocardiogram (ECG) in, 14-16
 pacemakers and, 16
 patient evaluation, 71-82
 testing, 72t-73t
 Diagnosis of thromboembolic stroke, 287
 DIAMOND trial, 29, 122, 212, 240-241
 Diaphragmatic paralysis, 172
 Dicloxacillin, 318t
 Diet, 320-321
 low-salt, 83
 DIG study, 29
 Digitalis, 29, 118t-119t, 126, 205, 220, 275
 Digoxin, 74, 261-264, 264t
 action mechanism, 261
 AF resulting from, 54
 clinical studies, 117, 126
 combination therapy, 265, 266, 275, 278
 contraindications/cautions, 154, 263-264
 dosing, 263, 264t
 drug interactions, 219-220, 261, 267
 exercise stress testing and, 77-78
 indications for, 261-263
 for rate control, 255, 261-264, 264t, 274-275, 278
 IV, 274-275
 Diltiazem, 205, 260, 262t, 274
 cautions for, 260
 clinical studies, 118t-119t, 127
 combination therapy, 265, 266, 278
 drug interactions, 261
 DIONYSOS study, 229
 Direct thrombin inhibitors, 304, 305, 322-334
 action mechanism, 322
 dabigatran, 323-334

Disopyramide, 206t, 246-248
 action mechanism, 246
 adverse effects, 247
 clinical studies, 118t-119t, 247-248
 contraindications, 247
 dosing, 206t, 246-247
 drug interactions, 247
 implanted device threshold effects, 199t
 in maintenance of sinus rhythm, 246
 monitoring, 202t-203t
 pharmacokinetics, 246
 safety, 199, 206t
Disulfiram, 319t
Dobutamine, 57
Dofetilide, 157t, 207t, 237-241
 action mechanism, 207t, 237
 adverse effects, 157t, 207t, 209
 clinical studies
 DIAMOND, 29, 122, 212, 240-241
 EMERALD, 240
 SAFFIRE-D, 157t, 240
 contraindications/cautions, 137t, 238
 dosing, 157t, 207t, 238
 drug interactions, 238-239
 implanted device threshold effects, 199t
 indications for, 157t, 237-241
 maintenance of NSR, 207t, 212, 237, 240-241
 initiation of therapy, 213, 215t, 239
 monitoring, 200-201, 202t-203t
 pharmacokinetics, 237
 safety of, 157t, 199, 200, 207t, 209-212, 238-240
 success rate for cardioversion, 157t
Domperidone, 198t
Dronedarone, 207t, 222-233
 action mechanism, 207t, 222-224
 adverse effects, 207t, 224, 229-232
 clinical studies, 225-233
 ADONIS, 226, 229
 ANDROMEDA, 227
 ATHENA, 227-229, 230-231
 DAFNE, 225-226
 DIONYSOS, 229
 ERATO, 227
 EURIDIS, 226, 229
 PALLAS, 232-233
 contraindications/cautions, 222, 224-225
 dosing, 207t, 224
 drug interactions, 225, 264
 implanted device threshold effects, 199t

Dronedarone (continued)
 indications for, 205-209, 210, 222-223
 maintenance of NSR, 205-209, 207t, 210, 212, 222-233
 rate control, 258
 initiation of therapy, 213, 215t, 222
 monitoring, 201, 202t-203t
 pharmacokinetics, 222-223
 safety of, 197, 198, 199-200, 207t, 212, 222
 PALLAS study, 232-233
 stroke risk, 229, 232
Droperidol, 198t
Drug interactions. See *specific agents*.
Dual-chamber defibrillators, 149-150
Dual-chamber pacemakers, 269, 270-271
Dyspnea, 68, 69
Echocardiography, 75-76, 203t, 204
 risk assessment with, 59-60
 stroke risk assessment with, 301-303, 303t
 transesophageal (TEE), 71, 73t
 transthoracic (TTE), 71, 72t, 75-76
Edoxaban, 334, 347-351
 bleeding risk, 350-351, 354t-355t
 clinical trials, 347-351, 350
 ENGAGE AF-TIMI 47, 347-351, 350
 in treatment algorithm, 305
EHRA scoring system, 73, 74t
Elderly patients. See also *Age*.
 amiodarone cautions for, 212
 case study, 391-393
 choice of therapy for, 124, 129, 129t
 degeneration of conduction system in, 270
 digoxin cautions for, 264
 hospitalizations in, 29, 30
 hypothyroidism in, 76
 lifetime risk and, 26
 neurologic evaluation, 78
 prevalence/incidence of AF in, 23-25, 24
 rate-control strategy in, 124, 129, 129t, 261, 274t
Electrical cardioversion, 134-150. See also *Cardioversion*.
 checklist for prior to, 134, 135t
 complications of, 142-143
 DC cardioversion, 134, 135t
 efficacy of, 133
 failure of, 143-144, 144, 146
 pharmacologic facilitation of, 147-149, 148t
 indications for, 133-134, 150-151
 in monitoring of AADs, 202t, 204

- Electrical cardioversion (*continued*)

outcomes and further management, 143-150

failed cardioversion, 143-144, 144, 146

pharmacologic facilitation of, 147-149, 148t

recurrence of AF, 143-144, 144

repeat cardioversion, 135, 145, 147, 147t

 AADs facilitating, 143, 143t

restoration of NSR, 143-144

practical and technical aspects, 135-139

anesthesia, 135-136

anticoagulation, 136-137

energy delivered, 139

high energy shock, 139, 147t

life-support capabilities, 135-136

pads/paddle use, 137, 138, 141

shock-induced arrhythmias, 149

shock synchronization, 137-139

shock waveforms, 139, 140

special populations, 141-142

 acute MI, 141

 hyperthyroidism, 141-142

 patients with implanted device, 141

 post-cardiac surgery, 141

 pregnancy, 142

success rate, 134

Electrical remodeling, 88, 150, 165

Electrocardiogram (ECG), 13-16, 15, 75

 in AAD monitoring, 201, 202t

ambulatory recordings/monitoring, 77, 79-80, 81t

of atrial flutter, 18

changes following cardioversion, 142

 δ wave on, 78

dilemma in diagnosis of AF, 14-16

of fibrillation, 15

follow-up monitoring, 78-80

in patient evaluation, 71, 72t, 75-76

risk assessment with, 59-60

Electrodes, placement for cardioversion, 137, 138, 141

Electrolyte testing, 77, 79, 80, 202t

Electrophysiology study (EPS), 71, 73t, 78, 79

Embolism. See *Thromboembolism*.

EMERALD trial, 240

Enalapril, 89-90

Endothelial and endocardial abnormalities, 283

Endurance sports exercise, 51, 52, 85, 123, 322

ENGAGE AF-TIMI 47, 347-351, 350

Epidemiology of AF, 23-32

 consequences and prognosis, 27-31

 demographics, 27

Epidemiology of AF (*continued*)

increase in AF, 23, 24

lifetime risk, 26

prevalence and incidence, 23-26, 24, 25

risk factors, 26

 EPOCH study, 52

ERAFT, 245

ERATO study, 227

Erythromycin, 335

ESC Guidelines. See *ACC/AHA/ESC Guidelines*.

Esmolol, 259t, 274

Ethnicity. See *Race/ethnicity*.

Etiology of AF, 47, 48t, 62. See also *Risk factors*.

Etomidate, 136

EURIDIS, 226, 229

Euro Heart Survey for AF, 299

EuroHeart Failure Survey, 29

European Heart Rhythm Association (EHRA) scoring system, 73, 74t

European Society of Cardiology. See *ACC/AHA/ESC Guidelines*.

Evaluation of patient with AF, 65, 71-82, 273

 follow-up encounters, 78-80

 minimum evaluation, 72t

 monitoring strategies, 81t

new-onset evaluation, 71-78, 72t-73t, 80

 blood tests, 76-77, 80

 chest imaging, 77

 echocardiogram, 75-76

 electrocardiogram, 72t, 75-76, 80

 electrophysiology study, 71, 73t, 78

 exercise stress testing, 72t, 77-78, 80

 history, 65, 71-74, 72t, 80

 physical examination, 65, 74-75

 symptom scoring systems, 73, 74t

Exercise, 85, 104

AF risk and, 52, 85, 86t-87t

exercise tolerance, treatment of AF and, 123, 123

stress testing, 71, 72t, 77-78, 80, 203t, 204, 277-278

walking, 85, 86t-87t

Extracellular matrix (ECM), 38-39, 85, 283, 285

F (fibrillatory) waves, 14

Factor Xa inhibitors, 304, 305, 334-351

 apixaban, 335, 339-347

 characteristics, 334-335

 edoxaban, 334, 347-351

 indications for, 334-335

 rivaroxaban, 335-339

Family history of AF, 47, 73

Fatigue, 68, 69

N-3 Fatty acids, 102
Fentanyl, 136
Fibrates, 318t

Fibrosis, 36, 39, 49, 85, 88, 212

First-detected episode, 19

Fish oil, 89, 102, 318t

Flecainide, 158t

adverse effects, 158t

clinical studies, 118t-119t, 125

dosing, 158t

drug interactions, 158t

implanted device threshold effects, 199t

indications for, 158t, 205-209, 206t, 210

pharmacologic cardioversion, 152-153

repeat electrical cardioversion, 143, 143t

monitoring, 202t-203t

safety, 198, 199, 200

success rate, 158t

Fluconazole, 318t

Fluorouracil, 319t

Fluvastatin, 319t

Focal activation, 33-36, 37, 165, 166

Foci/sites of initiation, 33-36, 37

Follow-up evaluation, 78-80

Fondaparinux, 322

Framingham risk score, 60, 61, 300, 302

Framingham Study, 23, 26, 28, 29, 52, 56, 60

Gender

AF incidence and prevalence and, 24, 25, 27

in CHA2DS2-VASc score, 298t

thromboembolic risk and, 292, 293t, 298t

Gene therapy, 273

Genetic AF, 49-51, 62

genetic etiology, evidence for, 47

genetic polymorphisms, 315-316

genetic variants, 34

susceptible phenotypes, 85

GI agents, 319t

GISSI-AF trial, 93

Glucocorticosteroids, 89, 90t, 98-101, 100, 104

safety of, 101

Goals of therapy, 109-114, 110

Griseofulvin, 318t

Growth factors, 88, 285

Haloperidol, 198t, 319t

HAS-BLED scoring system, 314-315

Heart disease

AF and, 27, 47, 53

statins and, 92

valvular, 53

Heart failure (HF), 28-29, 52-53, 61, 69

AAD therapy and, 194, 212

AF with, 122, 126-127, 271

AF-CHF trial, 122, 126-127

decompensated, 65, 275

pacemaker implantation, 271

Heart rate, 14, 16, 254, 258t. See also *Rate vs rhythm control*.

chronotropic incompetence, 257

with exercise, 258t

outpatient monitoring, 78

rapid, 41

with rate-control therapy, 254, 255, 258t

lower limit, 257, 258t

target (goal), 117, 254

upper limit, 258t

Heart sounds, 74

Hematocrit, elevated, 284

Hematuria, 314

Hemodynamic effects of AF, 28-29, 41-44

impaired coronary blood flow, 43

irregular ventricular rate, 41-43

loss of atrial contraction, 43

neurohormonal activation, 43-44

rapid heart rate, 41

Hemodynamic stability, achieving, 110

Hemodynamically unstable patients, rhythm control for, 137

HEMORR2HAGES risk index, 314, 315

Heparins, 362, 365

cardioversion and, 137, 171, 362

Herbal supplements, 320, 320t

Hispanics, 85

History, 65, 71-74, 72t, 80

HIV protease inhibitors, 198t

HMG-CoA Reductase Inhibitors (statins), 89, 90t, 93-98

Holter monitoring, 71, 73t, 81t

Hospitalizations for AF, 29-31, 30, 123, 127

in hospital telemetry, 81t

HOT CAFE trial, 116, 118t-119t, 122

HRS/EHRA/ECAS Consensus Statement, 174, 177, 295

Hydrocortisone, 99, 100

Hypercoagulable state, 28, 284-285

Hypertension, 51-52, 209
ACE inhibitors and, 92, 214
antihypertensive agents, 61, 305
in CHA2DS2-VASc score, 298t
in CHADS2 score, 294, 294t
clonidine for, 267-268, 278
treatment for, 61

Hyperthyroidism, 55-56, 141-142, 368-369
case study, 395-397

Hypertrophic cardiomyopathy (HCM), 54, 65, 68

Hypotension, 65, 74, 147

Hypothyroidism, 76

Ibutilide, 145t, 148-149, 156t, 164
implanted device threshold effects, 199t
indications for, 161
cardioversion of atrial flutter, 164
failed electrical cardioversion, 147-148, 148t
pharmacologic cardioversion, 148t, 156t
repeat electrical cardioversion, 143, 143t
monitoring, 149
safety of, 148-149, 156t, 161-164
torsade de pointes (TdP), 148, 155, 156t

Idiopathic AF, 47-51, 48t

IKur, 155

Imaging, 170, 287

Immediate recurrence of AF (IRAF), 144-146, 145t

Implanted devices. See also *Pacemakers*.
electrical cardioversion and, 141
interference from AADs, 199, 199t
threshold effects of AADs, 199t

Incidence of AF, 23, 24

Individualized approach to therapy, 109, 110, 111, 115, 126, 129, 129t, 203

Inflammation, 26, 36-38, 40, 48t, 60, 93. See also *Anti-inflammatory agents*.
AF prevention and, 89
anti-inflammatory agents, 40, 93, 98
thromboembolic risk and, 285

Initiating event, 33-36, 35

Inpatient settings (AADs), 152-153

Interferon, 318t

Internal cardioversion, 133, 134, 149

Ion currents, 160-161, 160

Irbesartan, 320

Isoniazid, 318t

Isoproterenol, 40

J-RHYTHM (Japanese Rhythm Management Trial for Atrial Fibrillation), 125-126

Joules, 139

Ketoconazole, 198t, 238-240, 324, 335

Laboratory data, 60

Lactation, 322

Left atrial appendage (LAA)
blood stasis and, 284
closure devices, 372-375, 373, 374
as source of thromboembolism, 286
surgical amputation/ligation, 381

Left atrial dilatation. See *Atrial dilatation*.

Left atrial pressure, elevated, 47, 48t

Left atrial vein ablation, 168, 169-170

Left atrium. See also *Atria*.
reservoir function, 38, 60
size/volume, 38, 60, 75

Left ventricular dilatation, 284

Left ventricular dysfunction, 28-29, 88

Left ventricular ejection fraction (LVEF), 41

Left ventricular end diastolic pressure (LVEDP), 43

Left ventricular hypertrophy (LVH), 52, 209

Life-support capabilities, 135-136

LIFE trial, 91

Lifestyle modifications, 83-85

Limb ischemia, 290t

Liver function tests, 72t, 77, 201, 202t

Liver toxicity, 200

LMWH, 137, 362, 363, 368, 370. See also *Heparins*.

Lone AF, 20, 47-49, 205
case study, 389-391

Long-standing persistent AF, 19

Loop diuretics, 198t

Loop recorders, 81t

Losartan, 91

Macrolides, 198t, 318t

Magnesium level assessment, 200, 202t

Magnesium sulfate, 268

Manitoba study, 29

Mechanism of AF, 33-40, 34
“AF begets AF,” 38-39, 85, 193
arrhythmia substrate, 36-38, 165
focal activation vs multiple wavelets, 36, 37
initiating event, 33-36, 35
modulating factors, 35, 40
multiple-source hypothesis, 36, 37
single-focus theory, 33-36, 37

Medical conditions associated with AF, 48t, 54-57
Medication history, 54
Medications, AF and, 57
Mefloquine, 198t
Megestrol, 238
Mesalamine, 318t
Mesenteric ischemia, 290t
Metabolic risk factors for thromboembolism, 293t
Metabolic syndrome, 55
Methadone, 198t
Methylprednisolone, 98, 99-101
Metoprolol, 259t, 274
Metronidazole, 317, 318t
Midazolam, 136
Milrinone, 57
Mitral stenosis, 284
 anticoagulation with, 297
Monitoring of AADs, 80, 200-204, 202t-203t
Monitoring of AF, 76-80
 recording options, 81t
Mortality, 27, 29, 115-116, 117-120
 catheter ablation and, 173, 173t
 hazard ratios for death in subgroups, 121
 stroke and, 285, 288
MOST, 103
MRI, 170, 287
Multifocal atrial tachycardia (MAT), 16, 54
Multiple-source hypothesis, 36, 37
Myocardial infarction (MI)
 AADs and, 115, 209
 acute, 53
 DC cardioversion and, 141, 151
 transient AF during, 58
 AF secondary to, 20, 47, 48t, 53
 precipitated by AF, 68
 transient AF and, 58
Myocardial necrosis, 142

N-acetylcysteine (NAC), 101
N-3 Polyunsaturated fatty acids (PUFAs), 102
NAC (*N*-acetylcysteine), 101
Nadolol, 258
Nafcillin, 318t
NASPE Registry, 271
NavX, 170
Neurohormonal activation, 43-44
Neurologic evaluation, 78
Niigata Preventive Medicine Study, 55
Nitrous oxide, 284

Non-valvular AF, 19
Nondihydropyridine CCBs. See *Calcium channel blockers*.
Nonpharmacologic therapy. See also *Implanted devices; Surgical ablation*.
 for prevention of AF, 83-85, 84
 for rate control, 269-273, 269t
 for stroke prevention, 372-381
Norepinephrine, 44
Normal sinus rhythm (NSR), 133-252. See also *Antiarrhythmic drugs (AADs)*.
 anticoagulation and, 134
 maintenance of, as a goal, 193
 AADs for, 193-252, 206t-207t, 210-211, 216-248
 indications for, 194
 restoration of, as a goal, 109, 115, 124, 129
 restoration and maintenance of, 133-252
 AADs for, 150-164, 193-200
 catheter ablation, 133, 151-153
 electrical cardioversion, 133, 134-150
 pharmacologic cardioversion, 150-164
 quality of life and, 122
 stroke and, 115, 121
 success rates, 144-145
NSAIDS, 317, 318t

Obesity, 55, 61
Obstructive sleep apnea, 55
Ocular effects, 219
Offset pause, 65, 69
Omeprazole, 371
ONTARGET trial, 92
Oral contraceptives, 319t
Organ toxicity, 200, 209, 212
Outpatient settings (for AADs), 152-153
Oxygen
 antioxidants, 101
 oxidative stress, 101
 reactive oxygen species, 88, 89, 93

P-glycoprotein inducers, 324
P waves, 13, 59-60
Pacemakers, 269-272, 279
 ACC/AHA/HRS device guidelines, 103-104
 AF diagnosis and, 16
 AF prevention and, 103-104
 choice/types of, 270-271
 dual-chamber, 269, 270-271
 physiologic pacing with, 104
 single-chamber, 269-270

Pads/paddles, placement for cardioversion, 137, 138, 141
 PAF/PAF-2 trials, 124-125
 PAI-1, 284
 Pain, electrical shock and, 136
 PALLAS study, 232-233
 Parasympathetic nervous system, 40, 48t
 Paroxysmal AF, 19, 38, 40, 165
 antithrombotic therapy in, 295
 clinical trials, 124-126
 and heart failure, 122
 rate vs rhythm control in, 128, 129t
 vs persistent AF, 167
 Pathophysiology of AF, 33-45, 34, 165. See also *Risk factors*.
 hemodynamic effects, 41-44
 impaired coronary blood flow, 43
 irregular ventricular rate, 41-43
 loss of atrial contraction, 43
 neurohormonal activation, 43-44
 rapid heart rate, 41
 mechanism of AF, 33-40, 34
 "AF begets AF," 38-39, 85, 193
 arrhythmia substrate, 36-38, 165
 inflammation and, 36-38, 48t
 initiating event, 33-36, 35
 modulating factors, 40
 Patient evaluation. See *Evaluation of patient with AF*.
 PAVE study, 271
 Pentamidine, 198t
 Percutaneous closure of LAA, 372-375, 373, 374
 Pericarditis, 48t, 53-54, 67t
 Peripheral thromboembolism, 28, 285, 288
 Permanent AF, 19
 Persistent AF, 19
 vs paroxysmal AF, 167
 Pharmacologic agents for rate control, 257-264
 Pharmacologic cardioversion, 150-164. See also *Cardioversion*.
 choice of AADs for, 154t
 new (atrial selective) AADs, 155-164
 indications for, 150-151
 pill-in-the-pocket strategy, 151, 158t, 213
 safety of, 151-153
 anticoagulation, 151
 inpatient vs outpatient settings, 152-153
 toxicity of AADs, 151-152
 success rate, 144-145, 150
 Pharmacologic prevention of AF, 85-102. See also *Prevention of AF*.
 modes of action, 90t
 Pharmacologic prevention of stroke in AF, 303-372
 Phenothiazines, 198t

Phenytoin, 319t
 Physical examination, 65, 74-75
 Physical exercise, 85, 86t-87t, 104, 123
 PIAF trial, 116, 118t-119t
 Pill-in-the-pocket strategy, 151, 158t, 213
 Pilsicainide, 125-126
 PLAATO, 372, 373
 Platelet activation, 284-285
 Platelet count, 76
 Platelet-derived growth factor (PDGF), 88
 Platelet factor 4, 284-285
 Pneumonia, 67t, 173t
 Polyunsaturated fatty acids (PUFAs), 102
 Postoperative AF, 48t, 57, 58-59, 59
 anticoagulation for, 368-369
 case study, 393-395
 fish oil and, 102
 pericarditis and, 53
 steroid therapy and, 98-101, 100, 104
 Potassium channels, 49, 207t
 Potassium level assessment, 200, 202t
 PR interval, 59, 61
 Prasugrel, 371
 Pravastatin, 95, 96t-97t
 Pre-excitation syndrome, 16, 57, 68, 275, 277
 Pregnancy
 DC cardioversion in, 142
 warfarin use in, 321-322
 PREVAIL trial, 377-378
 Prevalence of AF, 23-26, 25
 Prevention of AF, 83-108, 84
 device-based therapy in, 103-104
 goals of, 83, 84
 nonpharmacologic methods, 83-85, 112
 pharmacologic, 85-102, 89, 112
 AADs and, 214-215
 ACE inhibitors/ARBs, 88-93, 90t, 104
 aldosterone antagonists, 88, 90t, 93
 antioxidants, 101
 fish oil, 102
 glucocorticosteroids, 89, 90t, 98-101, 100, 104
 modes of action, 90t
 RAAS blockade, 88-93, 104
 statins, 90t, 93-98, 104
 target pathways, 89
 recurrence prevention, 205
 target pathways, 84, 89

Prevention of thromboembolism, 279-352, 283-388.
 See also *Thromboembolism, preventive therapies*.
 pharmacologic prevention in AF, 303-372, 305
 Proarrhythmias, 195-198
 atrial, 197-198
 ventricular, 195-197, 196, 212
 Procainamide, 148t, 154, 200
 Prochlorperazine, 238
 Prognosis of AF, 27-31
 Propafenone, 158t, 206t, 242-246
 action mechanism, 206t, 242, 267
 adverse effects, 206t, 243
 clinical studies, 118t-119t, 125, 244-245
 contraindications/cautions, 209, 212, 244
 dosing, 158t, 206t, 243
 drug interactions, 158t, 244, 261, 264, 318t
 implanted devices and, 199t
 indications for, 158t, 205-209, 210, 242-246
 maintenance of NSR, 206t, 242
 pharmacologic cardioversion, 158t
 pill-in-the-pocket approach, 151, 158t
 pretreatment for pharmacologic cardioversion, 148, 148t
 for rate control, 267
 repeat electrical cardioversion, 143, 143t, 146t
 monitoring, 202t-203t
 outpatient use of, 152
 pharmacokinetics, 242
 safety, 198, 199, 200
 Propofol, 136
 Propranolol, 274
 Protease inhibitors, 198t
 PROTECT-AF, 375-376, 377, 378
 Prothrombic state, 284-285
 Prothrombin time, 76, 313
 Prothrombotic plasminogen activator inhibitor 1 (PAI-1), 284
 PROVE IT-TIMI 22, 95
 Psychosocial factors, 26
 Psychotropic drugs, 198t
 PUFAs, 102
 Pulmonary disease, 54-55
 Pulmonary embolus (PE), 54, 67t
 chest CT for, 77
 Pulmonary vascular and pulmonary disease, 54-55
 Pulmonary vein isolation, 38, 168-169, 168
 Pulmonary vein stenosis, 171-172
 Pulmonary veins (PVs), in paroxysmal AF, 33-34
 Purple toe syndrome, 321

QRS complex, 75, 139, 201-204
 irregular, 14
 narrow, 14
 wide, 73t, 78
 QT intervals, 60, 75, 78
 prolonged, 60, 154, 155, 196-197, 196, 198t, 224
 congenital long-QT syndrome, 197
 drugs associated with, 198t, 213
 Quality of life (QoL), 27, 31, 109, 122, 124, 127
 Quinidine, 146t, 200, 324
 drug interactions, 264
 Quinolones, 198t, 318t
 R-R interval, 14, 75
 RAAS (renin-angiotensin-aldosterone system) activation, 34
 fibrosis and, 88
 RAAS (renin-angiotensin-aldosterone system) blockade, 88-93, 89, 104
 Race/ethnicity, 27, 52, 85
 RACE trial, 116, 118t-119t, 121, 122, 216, 254
 RACE II, 255-256
 Radiofrequency ablation, 187-188, 270
 Ramipril, 92
 Ranolazine, 198t
 Rate control, 115, 133-192, 253-282. See also *Heart rate*.
 in AF, 269-273
 algorithms based on clinical presentation, 273-279, 276-277, 280
 acute pharmacologic rate control, 273-275, 276-277
 chronic pharmacologic rate control, 277-279, 280
 decompensated heart failure (HF), 65, 275
 sick sinus syndrome or AV block, 277
 special populations, acute rate control in, 275-277
 Wolff-Parkinson-White (WPW) syndrome, 275-277
 bradycardia and, 149, 256-257
 caution in, 274-275, 274t
 choice of, vs rhythm control, 127-129, 129t
 clinical benefit of, 271
 clinical trials on, 116-124, 118t-119t
 pharmacologic agents used, 116-117
 combination therapy, 215, 275
 cost-effectiveness of treatment, 124
 effectiveness of, 127-128
 goals and definitions in, 253-257, 254t
 indications for, 128-129, 129t, 208, 253, 254
 acute presentation, 254t, 262t, 264t
 cautions, 274-275, 274t
 chronic presentation, 254t, 262t, 264t
 patient characteristics, 129t
 lenient, vs strict, 255-256

Rate control (*continued*)

nonpharmacologic therapy for, 269-273, 269
AV nodal ablation with pacemaker implantation, 270-272, 277, 279
AV nodal modification, 272
chronic atrioventricular nodal vagal stimulation, 273
gene therapy, 273
pacemaker implantation, 269-270
regularization of ventricular response with pacing, 272-273
pharmacologic agents for, 255, 257-264, 273-275
acute control with, 254t, 262t, 264t, 274, 275-277, 276-277
antiarrhythmic drugs, 258, 264, 266-267
 amiodarone, 258, 266-267
 propafenone, 267
 sotalol, 267
 β-blockers, 255, 258-260, 259t, 260t, 265, 273-274, 278
 cautions in titration of therapy, 274t
 CCBs, 255, 260-261, 265, 273-274, 278
 chronic control with, 254t, 262t, 264t, 277-279, 280
 combination therapy, 274, 275
 comparative studies, 265, 266
 digoxin, 255, 261-264, 264t, 278
 diltiazem, 260, 262t
 other agents
 clonidine, 267-268
 magnesium sulfate, 268
 verapamil, 260-261, 262t
 strategies for, 253-282
 target heart rate, 117, 254
 vs rhythm control. See also *Rate vs rhythm control*.
 therapy, 115-131, 118t-119t

Rate vs rhythm control, 115-131, 118t-119t, 253
age and, 124, 129t
choice of strategy, 112, 124, 127-129, 129t
clinical trials: agents used, 116-117, 118t-119t, 125-126
clinical trials: outcomes, 116-124, 118t-119t
 cost-effectiveness, 124
CV mortality, 120, 120
exercise tolerance, 123, 123
heart failure, impact on, 122
hospitalization, 123
symptoms and QoL, 31, 122
thromboembolic complications, 120-121
total mortality, 117-120
clinical trials: overview, 116-117, 118t-119t
clinical trials: specific populations, 124-127
AF and heart failure, 126-127
symptomatic, paroxysmal AF, 124-126
young patients with AF and valvular heart disease, 127

Rate vs rhythm control (*continued*)

goals of, 115, 128-129, 129t
 normal sinus rhythm, 115, 128-129
patient characteristics and, 129t
safety and efficacy, 115-116, 119t
stroke rates and, 292
thromboembolism risk, 292
Reactive oxygen species, 88, 89, 93
Recurrence of AF, 19, 173-174
 AADs in prevention of, 205, 213
 booster dosing, 213
catheter ablation and, 173-174
electrical cardioversion and, 143-146, 144
immediate recurrence (IRAF), 144-146, 145t
RE-LY trial, 325-330, 326, 327t
RELY-ABLE trial, 330-331
Remodeling. See *Atrial remodeling*.
Renal function tests, 72t, 77, 200, 202t
Renal thromboembolism, 291t
Renin-angiotensin-aldosterone system. See *RAAS*.
Retroperitoneal hemorrhage, 314
Rheumatic heart disease (RHD), 53, 127
Rhythm control, 115, 133-252. See also *Antiarrhythmic drugs*;
 Normal sinus rhythm; specific topics.
antiarrhythmic drugs (AADs) for, 193-252
anticoagulation therapy and, 215-216
catheter ablation, 164-184
 choice of, vs rate control, 127-129, 129t
 clinical trials on, 118t-119t
 electrical cardioversion, 134-150
 hospitalizations and, 127
 indications for, 126, 128-129, 129t, 133, 194, 275
 in hemodynamically unstable patients, 137
 patient characteristics, 129t
 maintenance of normal sinus rhythm, 193-252
pharmacologic agents for, 116-117
pharmacologic cardioversion, 150-164
quality of life and, 31
vs rate control, 115-131, 118t-119t. See also *Rate vs rhythm control*.
RIETE score, 314
Rifampin (rifampicin), 318t, 335
Right ventricular (RV) pacing, 103
RIKS-HIA, 263
Risk assessment, 59-60, 61
 for stroke, 289-303
Risk factors, 25-26, 47-64, 48t, 83
 and AF prevention, targets for, 84
assessment with ancillary studies, 59-60, 61
cardiac/noncardiac conditions associated with AF, 48t, 51-58, 62

Risk factors (*continued*)

alcohol and other toxins, 56-57
 cardiac surgery, 57, 58-59, 59
 congenital heart disease, 54
 coronary artery disease and acute MI, 53
 heart failure, 52-53
 hypertension, 51-52
 hyperthyroidism, 55-56
 hypertrophic cardiomyopathy, 54
 medications, 57
 metabolic syndrome, 55
 obesity, 55, 61
 pericarditis, 53-54
 pulmonary vascular and pulmonary disease, 54-55
 supraventricular arrhythmias, other, 57-58
 valvular heart disease, 53
 CV risk and, 85
 idiopathic AF, physiologic, and genetic factors, 47-51
 aging, 49, 61
 endurance sports exercise, 51, 52
 genetic AF, 49-51, 62
 lone AF, 47-49
 lifetime risk, 26
 prevalence of AF and, 26
 reversible risk factors, controversies on, 58-59
 risk score assessment/algorithm, 59-60, 61
 for stroke, 120-121, 289-303
 Risk score, 59-60, 61
 Risperidone, 198t
 Ritonavir, 198t, 335
 Rivaroxaban, 335-339
 action mechanism, 335
 as alternative to warfarin, 351
 clinical trials, 336-339
 ROCKET-AF, 336-339, 337, 338t
 FDA-approved uses, 334
 indications, risk category and, 296t
 indications for, 334-335
 in treatment algorithm, 305
 Rochester Epidemiologic Project, 23, 26
 ROCKET-AF, 336-339, 337, 338t
 Rotterdam Study, 23, 28, 57, 102
 SAFE-T, 122, 123, 123, 159t, 220, 235, 236
 SAFFIRE-D trial, 157t, 240
 SAVEPace trial, 103
 Scoring systems, 61, 73, 74t
 for thromboembolism risk, 294-300, 294t, 298t
 Secondary AF, 20

Sertraline, 319t
 Serum electrolytes, monitoring, 77, 79, 80
 Shock therapy. See *Electrical cardioversion*.
 Shock waveforms, 139, 140
 Sick sinus syndrome, 48t, 58, 258-260, 277
 Single-focus theory, 33
 Sinoatrial (SA) nodal cells, 13
 Sinus arrest, 142
 Sinus rhythm. See *Normal sinus rhythm (NSR)*.
 Skin effects, 219, 321
 Sleep disorders, 55, 73
 Smoking (tobacco), 57
 cessation of, 83
 Social history, 73
 Sodium channel blockade, 206t
 Sodium channel genes, 49
 SOLVD (Studies of Left Ventricular Dysfunction), 28-29, 89-90
 Sotalol, 207t, 233-236
 action mechanism, 207t, 233, 267
 adverse effects, 207t, 234-235
 clinical studies, 116, 118t-119t, 122, 125, 235-236, 236
 CTAf, 122, 221, 222, 235-236
 SAFE-T, 122, 123, 123, 159t, 220, 235, 236
 contraindications/cautions, 154, 212, 235
 dosing, 207t, 234
 initial, 214
 drug interactions, 235
 implanted device threshold effects, 199t
 indications for, 205-209, 210, 233-236
 maintenance of NSR, 205-209, 207t, 210, 212, 234, 235-236
 for rate control, 267
 repeat electrical cardioversion, 143, 143t
 initiation of therapy, 213-214, 215t
 monitoring, 200-201, 202t-203t, 214
 pharmacokinetics, 234
 safety, 198, 200, 207t, 209-212
 torsade de pointes, 197
 SPAF I trial, 76, 293, 304, 357-358
 SPAF III trial, 294, 307
 Special populations
 acute rate control in, rate control, 275-277
 electrical cardioversion in, 141-142
 thromboembolism prevention in, 359-372
 SPINAF trial, 287, 293, 306
 Splenic infarction, 291t
 Spontaneous cardioversion, 150
 Sports exercise, 51, 52, 85, 322
 SSRIs, 198t, 319t
 STAF trial, 116, 118t-119t, 122

Statins, 89, 93-98, 104
 actions of, 90t, 93
 combined with β -blockers, 95-98
 drug interactions, 318t
 for stroke prevention, in treatment algorithm, 305
 Stents, 397-399
 coronary stents, 371-372
 stent thrombosis, 369-370
 Steroids. See *Glucocorticosteroids*.
 STOP-2, 91
 STOP-AF, 176
 Stress (exercise) testing, 71, 72t, 77-78, 80, 203t, 204, 277-278
 Stroke, 285-288. See also *Thromboembolism*.
 age and, 286, 293t
 anticoagulation in patients presenting with, 364-365
 cardioembolic, 286-287, 288
 gender and, 293t
 ischemic, 286-288
 mortality and, 285, 288
 prevention in AF, 113, 279-352. See also *Thromboembolism*.
 algorithm for, 305
 anticoagulation therapy and, 120-121, 364-365
 clinical studies. See *SPAF I; SPAF III*.
 pharmacologic prevention, 303-372, 305
 prevention trials, 293-294
 prior
 in CHA2DS2-VASc score, 298t
 in CHADS2 score, 294, 294t
 risk, 292-297
 AF and, 27-28
 cardioversion and, 136
 in community patients with new-onset AF, 300-301, 302
 dronedarone and, 229, 232
 stratification schemes in AF, 292-297, 293t
 warfarin reduction of, 121, 129
 silent, 287
 Substrate for arrhythmia, 36-38, 84, 165
 Sucralfate, 319t
 Supraventricular arrhythmias, other, 57-58, 79
 Supraventricular tachycardia (SVT), 57
 Surgery, cardiac
 electrical cardioversion and, 141
 increase in AF after, 40, 57, 58-59, 59
 Surgical ablation, 184-187, 185, 270-272, 381
 AV nodal ablation with pacemaker implantation, 270-272, 277
 benefits vs risks, 186, 186t, 187, 188t
 catheter ablation and, 164-184
 complications of, 187

Surgical ablation (*continued*)
 Cox-Maze procedure, 184-187, 185, 186t
 Cox-Maze III, 186-187, 186t, 188
 Cox-Maze IV (bipolar radiofrequency), 187-188, 187
 benefits of, 188t
 risks, 185-186, 186t, 271-272
 success rates, 184, 184t
 techniques, 166-170, 168
 variations on, 184-185
 Sympathetic nervous system/tone, 40, 48t, 73, 89
 Sympathomimetic drugs, 57, 74
 Symptoms of AF, 13, 41, 65-70, 122. See also *Clinical manifestations*.
 asymptomatic AF, 31, 65, 69
 cardioversion and, 135
 hemodynamic, 41-44
 improvement of, as goal of therapy, 122
 palpitations/fluttering, 65
 poor correlation with AF frequency/duration, 68, 78
 relief of, as goal of therapy, 109
 scoring/classification of, 73, 74t
 severe, treatment choice and, 124, 129, 129t, 133, 135
 Syncope, 65-68, 69
 Systolic blood pressure, 61
 T wave, 137
 Tachycardia
 atrial, 16, 17, 17
 AV nodal reentrant, 17
 AV reentrant, 17
 supraventricular, 57
 ventricular, 16
 Tachycardia-induced cardiomyopathy, 28, 41, 42, 65, 69, 128
 Tamoxifen, 319t
 Tamponade, 172-173, 173t
 TEE. See *Transesophageal echocardiography*.
 Telmisartan, 92
 Terfenadine, 198t
 Tetracycline, 318t
 TGF- β , 88
 Theophylline, 54, 74
 Therapy. See also *Prevention of AF*.
 antiarrhythmic drugs, 193-252
 cardioversion, 133-192
 catheter ablation, 133, 164-184
 choice of strategy, 112, 124, 127-129, 129t
 device-based, for AF prevention, 103-104
 goals of, 109-114, 110
 acute/primary, 109, 110

- Therapy, goals of (*continued*)
 - long-term, 109, 110
 - individualized approach to, 109, 110, 111, 115, 126, 129, 129t, 205
 - nonpharmacologic
 - for AF prevention, 83-85, 112
 - for rate control, 269-273, 269t
 - for stroke prevention, 372-381
 - normal sinus rhythm, restoration of, 133-192
 - pharmacologic, for AF prevention, 85-102, 112
 - pharmacologic, for stroke prevention, 113
 - rate vs rhythm control, 112, 115-131, 118t-119t
 - rate control, 253-282
 - goals of, 253-257, 254t
 - nonpharmacologic therapy, 269-273, 269t
 - pharmacologic agents, 257-264
 - rhythm control, 133-252
 - antiarrhythmic drugs (AADs) for, 193-252
 - available agents, 216-248
 - catheter ablation, 133, 164-184
 - electrical cardioversion, 134-150
 - pharmacologic cardioversion, 150-164
 - surgical ablation, 184-187, 185
 - therapeutic approaches, 112-113
 - preventive therapy, 83-108, 112
 - stroke prevention, 113
 - antiplatelet therapy, 304, 357-359
 - nonpharmacologic methods, 372-381
 - surgical techniques, 381
 - surgical options, 184-187, 185, 381
 - Thiazide diuretics, 198t, 238
 - Thienopyridines, 369-370
 - Thioridazine, 198t
 - Thrombin inhibitors, direct, 304, 305, 322-334
 - Thromboembolism, 27-28, 283-388. See also *Stroke*.
 - AF and risk for, 27-28
 - anticoagulation therapy, duration of treatment with, 121, 124, 129, 134
 - cardioversion and, 136
 - catheter ablation and, 134
 - clinical presentation in AF, 285-288
 - cardioembolic strokes, 286-287
 - cardioversion and, 285
 - cerebral thromboembolism, 286
 - diagnosis, 287
 - immediate recognition and treatment, 285
 - ischemic stroke and TIAs, 286-288
 - other complications, 288, 290t-291t
 - outcome and mortality, 120-121, 288
 - complications, 120-121, 143
 - Thromboembolism (*continued*)
 - CVA, 65, 69, 172
 - diabetes and, 20
 - low-risk patients, 297, 300
 - pathophysiology in AF, 283-285
 - blood stasis, 284
 - endothelial and endocardial abnormalities, 283
 - hypercoagulable state, 28, 284-285
 - peripheral, 28, 285, 288
 - preventive therapies, 113, 279-352, 283-388, 296t
 - ACC/AHA/HRS Guidelines (2014), 289, 295
 - ACCP guidelines (2014), 295, 296t
 - agents available
 - aspirin, 296t, 357-358
 - dabigatran, 296t, 305, 323-334
 - warfarin, 296t, 304-322
 - anticoagulation therapy, 129, 304-372, 305
 - warfarin, 304-322
 - antiplatelet therapy, 304, 357-359
 - bleeding risk and, 306
 - LAA amputation or ligation, 381
 - newer anticoagulants, 304, 322-357
 - dabigatran etexilate, 323-334
 - factor Xa inhibitors, 304, 334-351
 - nonpharmacologic methods, 372-381
 - LAA amputation or ligation, 381
 - LAA closure devices, 372-375, 373, 374
 - in patients with AF, 296t
 - pharmacologic stroke prevention, 303-372, 305
 - basic strategies, 303-304, 305
 - special situations, 359-372
 - AF patients presenting with stroke, 364-365
 - AF patients with coronary interventions, 369-372
 - atrial flutter, 365-368
 - cardioversion of AF, 359-363
 - treatment algorithm, 305
 - warfarin, 296t, 304-322
 - alternatives to, 295
 - prothrombotic state, 284-285
 - in rate- vs rhythm-control trials, 120-121
 - risk stratification in AF, 289-303
 - age and, 286, 292, 293t
 - Birmingham scheme, 298, 298t, 300, 309
 - CHA2DS2-VASc, 298-300, 298t
 - CHADS2 score, 294-297, 294t, 300
 - clinical risk factors, 292-297, 293t
 - Framingham risk score, 300, 302
 - gender and, 293t, 298t
 - limitations of, 289

Thromboembolism, risk stratification in AF (*continued*)

low-risk patients, 297, 300
 in new-onset AF, 300-301, 302
 preventive therapies and, 296t
 rhythm- vs rate-control, 292
 scoring systems, 294-300, 294t, 298t
 stroke prevention trials, 293-294
 stroke risk based on echocardiographic assessment, 301-303, 303t
 stroke risk in community patients with new-onset AF, 300-301, 302
 Swedish Cohort Atrial Fibrillation Study, 293t
 TTE assessment for, 76
 type of AF, 292
 valvular vs nonvalvular AF, 292
 surgical ablation and, 381
 Virchow's triad of thrombogenesis, 285
 Thromboglobulin, 284-285
 Thyroid conditions: hyperthyroidism, 55-56, 141-142
 Thyroid hormones
 antithyroid drugs, 319t
 testing, 56, 60, 72t, 76, 79, 201
 Thyrotoxicosis, 258
 TIA, 286-288, 365
 Ticagrelor, 371
 Toes, purple/blue toe syndrome, 321
 Torsade de pointes (TdP), 148
 AADs and, 148, 155, 195, 196-197, 196, 213, 238
 TRACE study, 89
 Trandolapril, 89
 TRANSCEND study, 92
 Transesophageal echocardiography (TEE), 71, 73t, 361-363
 blood stasis on, 284
 prior to cardioversion, 80, 136-137
 prior to catheter ablation, 170, 171
 stroke diagnosis with, 287
 stroke risk assessment with, 301-303, 303t
 TEE-guided cardioversion, 361-363
 Transforming growth factor- β , 88, 285
 Transient ischemic attacks (TIAs), 286-288, 365
 Transthoracic echocardiography (TTE), 71, 72t, 75-76, 80
 repeat, 78-79
 stroke risk assessment with, 287, 303t
 Transthoracic procedure for cardioversion, 134
 Treatment of AF. See *Therapy*.
 Tricyclic antidepressants, 198t
 Triggers of AF, 17, 33-36, 167
 prevention strategies and, 84
 Trimethoprim, 238
 Trimethoprim-sulfamethoxazole, 317, 318t

Triple therapy, 369-370, 371-372

TTE. See *Transthoracic echocardiography*.

UFH, 368, 370. See also *Heparins*.

Vagal predominance, 40
 Vagal stimulation, 273
 Vagal tone, 57, 74
 Val-HeFT, 91
 Valsartan, 91
 VALUE trial, 91
 Valvular heart disease, 53, 127, 292
 AF secondary to, 47
 Vaughan-Williams classification of AADs, 206t-207t
 Ventricular arrhythmias, 149
 Ventricular fibrillation, 139
 Ventricular proarrhythmias, 195-197, 196, 212
 Ventricular rate, 14, 16, 128
 goals, with rate-control therapy, 128
 regularization with pacing, 272-273
 Ventricular tachycardia, 16
 Verapamil, 260-261, 262t, 278
 AF recurrence and, 205
 clinical studies, 118t-119t
 contraindications/cautions, 260-261, 324
 drug interactions, 238, 261, 264
 for rate control, 260, 262t, 274, 278
 for repeat cardioversion, 143, 143t, 146t
 Vernakalant, 160-164, 164t
 action mechanism, 160-161
 approval in EU, 163
 dosing, 164t
 oral formulation, under study, 163
 safety, 162-163
 safety and efficacy, 161-164
 side effects, 162-163, 164t
 success rate, 162, 162, 164t
 use for cardioversion, 160-164, 164t
 Veterans Affairs study, 221
 Virchow's triad, 285
 Vitamin C, 89, 101
 Vitamin E, 101
 Vitamin K, 312-313, 317, 319t, 320
 Vitamin K antagonists (warfarin), 303. See also *Warfarin*.
 in treatment algorithm, 305
 warfarin, 304-322
 VKORC1 gene polymorphism, 316
 von Willebrand factor (vWF), 283, 285
 Voriconazole, 198t

Walking, 85, 86t-87t, 104
six-minute walk test, 72t, 255, 278
Warfarin, 304-322
action mechanism, 311
advantages and disadvantages of, 356t
alternatives to, 295, 351
bleeding risk and, 306, 313-315, 317, 326-329, 328t
scoring systems, 314-315
stroke prevention and, 315, 316
catheter ablation and, 171
characteristics of, 310-322
clinical trials, 304-306, 325-331
contraindications, 372
dosing, 312-313
duration of treatment with, 121, 124, 129
efficacy
of low-intensity warfarin, 307-310
in primary prevention of stroke in AF, 304-307
in secondary prevention of stroke in AF, 307
indications for
age and, 309
based on CHADS2 and CHA2DS2-VASc scores, 297, 298
mitral stenosis, patients with, 297
prior to cardioversion procedures, 136
prior to catheter ablation, 171
interactions, 315-321
cholesterol embolization, 321
diet, 320-321
drug interactions, 317-320, 318t-319t
genetic polymorphism, 315-316
herbal dietary supplements, 320, 320t
hypermetabolism, 321
pregnancy/lactation and, 321-322
skin necrosis, 321
sports activities, 322
interruption of therapy, 370, 371
limitations of, 309-310
patient compliance and, 310
in rate vs rhythm trials, 117, 120
for stroke prevention, 120, 296t, 304-322
action mechanism, 311
bleeding risk and, 313-315, 316, 317
characteristics of, 210-222
dosing, 312-313
monitoring, 313, 314t
pharmacokinetics, 311-312
risk category and, 296t
vitamin K levels and, 312-313
vs dabigatran, 325-331

Warfarin (*continued*)
vs edoxaban, 347-351, 350
vs rivaroxaban, 336-339, 337, 338t
Watchman device, 372, 373, 374, 375-379
Weight loss, 83
WOEST trial, 371
Wolff-Parkinson-White (WPW) syndrome, 58, 68, 263-264, 275-277
Women's Health Study, 56-57, 101, 102
WPW syndrome. See *Wolff-Parkinson-White syndrome*.

X-ray, chest, 71, 73t, 77
Ximelagatran, 323

Young patients, 127, 128, 129, 129t, 133