

INDEX

Note: HF stands for heart failure.

Page numbers in *italics* indicate figures.

Page numbers followed by a “t” indicate tables.

Clinical trials and studies are indexed under the acronym of the name.

6-minute walk test, 40, 195, 196, 200, 226, 242, 249

A-type natriuretic peptide (ANP), 51, 88

ACC/AHA

guidelines for heart failure management, xvi-xviii[t], 258

stages of heart failure, 40-43, 41t

relationship to NYHA classification, 42

ACE inhibitors. See *ACEi-s*.

ACEi-s (angiotensin-converting enzyme (ACE) inhibitors), 83-86

action mechanisms, 83, 84

adverse effects, 86t

ARBs vs, 83-84, 87-88

cautions/contraindications, xv, xviii, 86t

clinical trials, 87-88

CHARM, 87-88

CONSENSUS, 84

ONTARGET, 87

SOLVD, 84

efficacy, 84, 87

indications for

HF-rEF patients, xviii, 84

HF with CKD, 194t

kinins and, 84

switching to an ARB, 86-87

switching to sacubitril/valsartan, 91, 94

Acute coronary syndrome (ACS), 128

Acute decompensated heart failure (ADHF), 253-277

classification of patients, 255, 258

cold and dry, 39t, 255, 258

cold and wet, 258

cold, cardiac output in, 257t

cold, treatment decisions for, 255

hemodynamic patterns, 39t

Acute decompensated heart failure (ADHF), classification of patients (*continued*)
 pulmonary congestion (“wet”), 256t
 treatment decisions and, 255
 volume status for “wet” vs “dry”, 32t, 259
 warm and dry, 255, 258
 warm and wet, 39t, 258
 clinical presentation, 254-255, 256t-257t
 low cardiac output (“cold”), 257t
 overall circulatory assessment, 255, 258
 pulmonary or systemic congestion (“wet”), 256t
 decompensation as sentinel prognostic event, 253
 goals of therapy, 255
 hemodynamic deterioration
 as marker for high risk of adverse outcomes, 264-266
 number of hospitalizations, mortality risk and, 253-254,
 254
 preceding clinical events, 253
 heterogeneity in ADHF population, 253, 274
 in HF-pEF patients, 253
 in HF-rEF patients, 253
 hospitalization for, 34, 34, 253-254
 EVEREST study, 254
 mortality risk and, 253-254, 254
 number of HF hospitalizations and, 254, 254
 initial evaluation, 255, 258t
 medical management
 challenges in, 274-275
 diuretics, 259-260
 goals of therapy, 255
 heterogeneity in ADHF population, 253, 274
 individualization of, 255, 274
 initial evaluation and, 255, 258
 summary and evaluation, 274-275
 symptomatic relief, 258-274. See also *specific agents listed under symptomatic relief, below*.
 new-onset, 253
 population characteristics, 253
 precipitating factors, 255, 259t
 prevalence of, 253
 signs and symptoms, 254-255, 256t-257t

Acute decompensated heart failure (ADHF) (*continued*)
 symptomatic relief, agents for, 258-274
 diuretics, 259-260, 261, 274
 dobutamine, 267t
 dopamine, 266-274, 267t
 vs nesiritide, 266-274, 270-273
 inotropes, 266-274, 267t
 not recommended for HF-rEF patients with ADHF, 266
 milrinone, 266, 267t
 adverse events with, 266, 268t-269t
 morphine, 258-259
 nesiritide, 264, 265t
 vs dopamine, 266-274, 270-273
 summary and evaluation of, 274-275
 ultrafiltration, 262, 274
 vasodilators, 262-266, 274
 blood pressure and, 264, 265t
 vasopressin antagonists, 260-262, 263
 Acute heart failure, 22t, 44t. See also *Acute decompensated heart failure (ADHF)*.
 Acutely decompensated HF (ADHF), 34, 39-40
 ADHF. See *Acute decompensated heart failure*.
 Adrenergic agonists, 266-274, 267t
 Adrenergic blockers. See β -blockers.
 Advanced HF, INTERMACS registry of stages for, 242,
 244t-245t, 248
 Afterload, 50
 Air pollution, 79
 Albuminuria, 61, 186
 Alcohol intake, 74-76, 75
 Aldactone. See *Spironolactone*.
 Aldosterone, 102
 Aldosterone antagonists, 102. See also *Eplerenone; Spironolactone*.
 agents, 102
 eplerenone, 102
 spironolactone, 102
 Alogliptin, 165t
 Alpha glucosidase inhibitors, 165t
 Altitude, 78
 Alzheimer’s disease, comorbid with HF, 160t
 American College of Cardiology/American Heart Association. See *ACC/AHA*.

- Amiodarone, 143
- Anemia, 66, 193-195
comorbid with HF, 160t, 162t, 193-195
iron deficiency, 193, 201, 202
iron supplementation treatments, 194-195, 196-200
- Angina, 36t, 46, 171
- Angioedema
as ACE-i side effect, 86t
contraindication for sacubitril/valsartan, xv, 89
minimizing risk for, xviii
prior, as medication contraindication, 86t
- Angiotensin, 50, 183. See also *Renin-angiotensin-aldosterone system*.
- Angiotensin I, 85
conversion to angiotensin II, 85
- Angiotensin II, 85
- Angiotensin II receptor blockers. See *ARBs*.
- Angiotensin-converting enzyme (ACE) inhibitors.
See *ACEi-s*.
- Angiotensin receptor neprilysin inhibitor (ARNI), xv, 88.
See also *Sacubitril/valsartan*.
- Anti-platelet agents, 125-127
- Anti-thrombotic agents, 125-131
- Anti-Xa anticoagulants, 127-128
- Antibiotics, 154-155
- Anticoagulants, 127-131
anti-Xa anticoagulants, 127-128
clinical trials
ATLAS-2, 131
COMMANDER-HF, 130-131
HF and AF, 127-128
HF with AF and CAD, 128-130
HF without AF, 130-131
implantable devices and, 152, 153
indications for, 127-128
oral, use with clopidogrel, 130
triple therapy, 129
warfarin, 127, 128
- Antidepressants, xx, 287
SSRIs, xix-xx, 287
tricyclic, xx
- Aortic stenosis, 217-219, 218
approved TAVR systems, 218-219
decision making for TAVR patients, 224
Edwards Sapien transcatheter valve system, 218
Medtronic Evolut R System, 218
post-procedure improvements, 220-223
- ARBs (angiotensin II receptor blockers), 83-84, 86-88
ACEi-s vs, 83-84, 88
action mechanisms, 83-84
adverse effects, 86, 86t
ARBs, 83-84, 86-88
cautions/contraindications, 86t
combining sacubitril/valsartan with, xv, xviii
clinical trials, PARAMOUNT trial, 60-61
discontinuation of, 87
dosage of, switching from ACEi to, 86-87
efficacy, 87-88
indications for, xix, 86-88
HF-pEF, 88
HF-rEF, 88
HF with CKD, 194t
switching to sacubitril/valsartan, 91, 94
tolerability, 86
- Arginine vasopressin, 260
vasopressin antagonists, 260-262, 263
- ARNI (angiotensin receptor neprilysin inhibitor), xv, 88.
See also *Sacubitril/valsartan*.
- PARAMOUNT trial, 60-61
- Arrhythmia. See *Atrial fibrillation*.
- Arthritis, comorbid with HF, 160t
- ASCEND-HF, 264, 265t, 274
- Aspirin, 129
- Asthma, 160t, 162t
- ATLAS-2 trial, 131
- Atrial fibrillation (AF), 26t, 127-131
ADHF precipitated by, 259t
- anticoagulants. See also *Anticoagulants*.
contraindication for, 228-229, 231
Watchman Device as alternative, 229-231, 230
- cardiac resynchronization therapy (CRT) for, 147-151
- cognitive impairment and, 282-283, 284-285
- heart failure plus CAD with, 128-130

- Atrial fibrillation (AF) (*continued*)
 heart failure with, 127-128, 160t
 heart failure without AF, 130-131
 incidence in HF patients, 228
 management of, left atrial appendage (LAA) closure, 228-231, 230
 prevalence, 127, 128
 progression of HF and, 35
 stroke risk, 127
 Atrium, left. See *Left atrial appendage (LAA) closure*.
 AV nodal ablation, 147-151, 151
 Avandia (rosiglitazone)
 Azotemia, 86t
- B-type natriuretic peptide (BNP), 51, 88
 increased, 256t
 using to evaluate HF with COPD, 169, 179
- β -Blockers, 96-99
 benefits of, xv
 efficacy, 96-98
 comparison of β -blockers, 98t
 differences between β -blockers, 97, 98
 in HF-pEF, 98-99
 vs ivabradine, 99
 indications for, 96, 98-99
 HF patients with COPD, 169-173, 181t, 183t
 in patients with HF-rEF, xviii
 ivabradine and, xv
- Bile acid sequestrants, 165t
 Biomarkers, 25t
 Biopsy, endomyocardial, 31
 Bisoprolol
 efficacy, 96
 indications for, HF with CKD, 194t
 Blood glucose, 25t
 Blood pressure. See also *Hypertension; Hypotension*.
 cognitive dysfunction and, 283
 vasodilators and, 264, 265t
 Blood urea nitrogen (BUN), 25t
 BMI (body mass index), 23, 73
 Body mass index (BMI), 23, 73
 Bradykinin, 84
- Brain natriuretic peptide, diagnostic importance of, 25t
 Brain natriuretic peptide. See *B-type natriuretic peptide (BNP)*.
 Bromocriptine mesylate, 165t
 Bumetanide, 115
 BUN (blood urea nitrogen), 25t
- CABG surgery, 159-161
 Caloric restriction, 73-74
 Canagliflozin, 165t
 Cancer, 202-210
 anti-HER2 and VEGF cardiotoxicity, 203, 210t
 anticancer drugs and their CV effects, 206t-207t
 cardiotoxic effects of anticancer agents, 202-203, 204-205
 CV risk assessment, monitoring, and management recommendations, 212t-213t
 frequency, mechanism, and reversibility of cancer therapies, 208t-209t
 QT-prolongation, 203
 radiation therapy, 203-210
 signaling inhibitor agents, 203, 207t
 Candesartan, clinical trials of, 87-88, 186, 190-193, 192
 Canrenone, 106-107
 Cardiac catheterization. See *Catheterization*.
 Cardiac decompensation. See *Acute decompensated heart failure*.
 Cardiac output, 39t, 64, 111
 assessment of, 31
 in cardiogenic shock, 39
 decreased, cognitive dysfunction and, 283
 in HF-pEF, 57
 in HF-rEF, 54
 low, 257t
 preload/afterload in, 50
 renal impairment and, 56t
 Cardiac resynchronization therapy (CRT), 143, 146, 148-149. See also *Pacing*.
 in AF patients, 147-151
 Cardiac surgery. See *Surgical treatments*.
 Cardiac transplantation, 30, 31, 73, 237
 Cardio-oncology, 202-210. See also *Cancer*.
 Cardio-renal syndrome (CRS), 55, 55t

- Cardiogenic shock, 22t, 39t, 244t
 hemodynamic patterns in, 39t, 244t
 inotropes for, 274
- CardioMEMS HF System, 236-237
- Cardiopulmonary function testing, 29-30, 30t
- Cardiovascular history, 23, 23t
- Cardiovascular implantable electrophysiologic devices (CIEDs), 141-158. See also *Implantable hemodynamic monitoring*.
 anticoagulation, 152
 cardiac resynchronization therapy (CRT), 143, 146, 148-149
 in patients with AF, 147-151, 151
 CIED therapy and clinical management, 152
 dyssynchrony induced by RV pacing, 147, 150, 150t
 implantation of a new device, 155
 importance and summary, 153-156
 infection, 152-153
 treatment of, 154-155
 QRS widening, 143, 146
 sudden death, 141-143
- Cardioversion defibrillators, implanted, 141-143, 144-145
 ACC/AHA guidelines, 143
- CARESS trial, 262, 274
- Carvedilol, 96-97
 efficacy, 96
 indications for, HF with CKD, 194t
- Catheterization
 ESCAPE trial, 39
 left heart, for diagnosis, 30-31
 of right heart
 clinical outcomes and, 39
 diagnostic, 31
- CHAMPION trial, 236-237, 238-241
- CHARM, 87-88, 186, 190-193, 192
- Chemotherapy, 23t
 cardiotoxic effects, 202-203, 204-205
- Chest X-ray, 27-28
 information from, 27t
- Chlorpropamide, 165t
- Chlorthalidone, 113-115
 combined with MRA, 115
 compared with hydrochlorothiazide, 114t
- Chronic heart failure, 22t, 34, 44t. See also *Nonpharmacologic management of chronic heart failure*; *Pharmacologic management of heart failure*.
 acute exacerbation of, 35
 acute exacerbation, vs slowly progressing, 35
 nonpharmacologic management of, 71-82
 pharmacologic management of, 83-140
 Chronic kidney disease (CKD), 185-193
 ADHF precipitated by, 259t
 comorbid with HF, 159, 160t, 185-193
 impact for HF-pEF, 190-193, 191, 192
 impact for HF-rEF, 186, 187t, 189-190, 192, 193
 Kaplan-Meier plots, 186, 188, 191, 192
 management principles, 193, 194t
 negative feedback cycle in, 186, 189
 prevalence and severity of hospital admissions, 189, 190
 prevalence of coexisting CKD in HF population, 186-189, 190
 definition and pathophysiology, 185-186
 GFR in, 185, 186
 MDRD formula, 185, 186
 prevalence in general population, 187t
 prevalence of in HF population, 186-189, 190
 in HF-rEF patients, 187t
 Chronic obstructive pulmonary disease (COPD), 35, 60, 66, 169-185
 comorbid with HF, 159, 160t, 162t, 169-185
 β-blockers for, 169-173, 181t, 183t
 ivabradine for heart rate reduction, 173-179, 184t, 185
 recommended therapies, 173, 180
 using B-type natriuretic peptide levels to evaluate, 169, 179
 heart failure and, 37t, 60, 169-185
 HF-pEF and, 60, 66
 HF-rEF and, 173-179
 prevalence in HF population, 169
 SHIFT trial, 173-175, 182t-184t
 Chronic pulmonary hypertension, 60
- Chronic thromboembolic pulmonary hypertension (CTEPH), 38
- Chronotypic incompetence, 54, 68t, 99
- CIEDs. See *Cardiovascular implantable electrophysiologic devices*.

Circulatory support, mechanical, 237-246
Classification of heart failure, 33-47
ACC/AHA stages of heart failure, 40-43, 41t, 42
acute HF, 34, 44t
 acutely decompensated HF (ADHF), 34, 39-40, 255, 258
based on anatomy and function, 35-40, 38, 44t
 hemodynamic patterns, 39-40, 39t
HF-rEF, 38-39
left heart failure, 38-39, 38
right ventricular failure, 35-38, 38
based on timing of symptoms, 34, 44t
 acute vs chronic, 34, 44t
chronic HF, 34, 44t
 acute exacerbation, vs slowly progressing, 35
comorbid processes and, 35
demographic data, 36t-37t
heart failure syndromes, classification of, 35-39
NYHA classification, 40-43, 40, 41t, 42
phenotypes of HF patients, 43, 46-47
treatment context provided by, 43, 44t
Clinical syndromes and staging of HF, 33-47. See also *Classification of heart failure*.
Clinical trials. See *specific agents, treatments, and specific trials by name*.
Clopidogrel, 130
Coenzyme Q10, 76
Cognitive impairment, 282-283, 284-285
 atrial fibrillation and, 282-283, 284-285
dementia, 160t, 282
ESC Guideline recommendations on, 281t
silent cerebral ischemia (SCI), 282, 283
Colesevelam, 165t
COMMANDER-HF, 130-131
Comorbid conditions, 36t-37t, 49, 159-216. See also *specific conditions, eg, Diabetes mellitus*.
Alzheimer's disease/dementia, 160t, 282
anemia, 160t, 162t, 193-195
arthritis, 160t
asthma, 160t, 162t
atrial fibrillation, 127-128, 160t
cancer, 202-210
cardiovascular conditions and risk factors, 36t

Comorbid conditions (*continued*)
challenges to HF management, 159
chronic kidney disease (CKD), 159, 160t, 185-193
commonly associated with HF, 159, 160t, 162t-163t
COPD, 159, 160t, 162t, 169-185
coronary artery disease (CAD), 159-161
depression, 160t, 283-288
diabetes, 159, 160t, 161-169, 162t, 165t
geriatric conditions, 37t
HF and atrial fibrillation, 127-128
 with CAD, 128-130
HF-pEF and, 49, 66-67
 COPD, 60
hyperlipidemia, 160t
hypertension, 36t, 160t
ischemic heart disease, 160t
obesity, 162t
other medical conditions, 37t
progression of HF and, 35
psychosocial conditions, 37t
COMPASS-HF trial, 231-236
Computed tomography (CT)
CONFIRM-HF study, 195, 202
Congestion, pulmonary or systemic, 255, 256t, 258
CONSENSUS trial, 83
Contrast ventriculography, 30
COPD. See *Chronic obstructive pulmonary disease*.
Cordarone (amiodarone), 143
Corlanor. See *Ivabradine*.
Coronary angiography, 30-31
Coronary artery disease (CAD), 159-161
 CABG surgery, 159-161
 heart failure and, 36t, 159-161
 heart failure and AF with, 129-130
Coronary disease, progression of, 35
Cough, ACE inhibitors and, 86t
CTEPH (chronic thromboembolic pulmonary hypertension), 38
Dapagliflozin, 165t
Decompensated heart failure. See *Acute decompensated heart failure (ADHF)*.

Decompensation, as sentinel prognostic event, 253
Defibrillators: implantable cardioverter-defibrillators, 141-143, *144-145*
Depression, xix-xx, 283-288
adherence to therapy and, xix
antidepressants, xx, 287
clinical trials on, 287
cognitive behavioral therapy, xix
ESC Guideline recommendations on, 281t
evaluation instruments, 287
exercise and, 287-288
heart failure and, 37t, 160t, 283
HF-rEF, 287-288
HF prognosis and, 283, 286
hospitalizations and mortality and, 283-287, 286
prevalence of, 283
screening for, xix
SSRIs for, xix-xx, 287
worsening of symptoms, 287
Diabetes mellitus, 23t, 66, 161-169
comorbid with HF, 159, 160t, 161-169, 162t
effects of anti-diabetic agents, 161-169
optimal management strategy, 169
potential adverse effects of anti-diabetic medications in HF, 161-169, *167, 176-177*
heart failure and, 36t, 161-169
heart failure treatment in, 159
management of, 161-169, 165t
DPP-4 inhibitors, 165-166, 165t
efficacy data, review of, *170-172*
metformin, 165t, 166
SGLT2 inhibitors, 165t, 167-169, *168*
adverse effects, 169, *176-177*
CV outcomes trials, 178t
type 1 diabetes, 161
type 2 diabetes, 161-169
drugs used in treatment of, 161-169, 165t
efficacy data, review of, *170-172*
Diagnosis of heart failure, 21-32, 32t, 56. See also *Classification of heart failure: Clinical syndromes and staging*
cardiac imaging studies, 28-29, 29t
2D and Doppler echocardiography, 28, 28t

Diagnosis of heart failure, cardiac imaging studies (*continued*)
comorbidities and, 29
important data from, 28t
chest X-ray, 27-28, 27t
electrocardiogram (12-lead resting), 24-27, 26t
exercise testing in, 30, 30t
functional studies, 29-30, 30t
history, 22-23
cardiovascular history, 23, 23t
initial evaluation in ADHF patient, 255, 258t
invasive studies, 30-31
cardiac catheterization, 30-31
contrast ventriculography, 30
coronary angiography, 30-31
endomyocardial biopsy, 31
right heart catheterization, 31
laboratory tests, 23-24, 25t
natriuretic peptides in, 25t
outcomes of HF evaluation, 32t
physical examination, 23, 24t
presentation, 21, 22t, 43, 256t-257t
salt and water retention, exercise intolerance, and dyspnea, 21, 22t
Diet, 71-77
alcohol intake, 74-76, 75
caloric restriction, 73-74
dietary supplements, 76-77
coenzyme Q10, 76
nitrate supplementation, 76-77
fluid restriction, 72
sodium (salt) limitation, 71-72
clinical evidence for, 71
conversion factors, 72t
RAAS activation associated with, 71
summary of recommendations, 74t
DIG trial, 116-117, *126, 190, 191*
Digoxin, 116-117, *126*
adverse effects, 116
clinical trials of (DIG trial), 116-117, *126*
indications for, 117
monitoring of serum digoxin levels, 117
risk ratio, 116-117

- Diuretics, 113-116. See also *specific agents*.
 adverse effects, 115
 agents
 chlorthalidone, 113-115, 114t
 furosemide, 33, 115, 122t-124t
 hydrochlorothiazide, 113, 114t
 torsemide, 115-116, 118-119, 120-121, 125t
 clinical trials, 113-115
 ALLHAT, 113
 DOSE, 259-260, 261, 274
 ROSE trial, 266-274, 270-273, 274
 combined with MRA, 115
 comparison of, 114t, 122t-124t, 125t
 continuous infusion with, 274
 dosage, low-dose vs. high-dose strategy in ADHF,
 259-260, 261
 efficacy, 113-116
 indications for, 113, 116
 in acute decompensated heart failure, 259-260, 261
 loop, 115-116
 thiazide, 113-115
 Dobutamine, 267t
 Dopamine, 266-274, 267t
 low-dose (“renal-dose”), to enhance effects of diuretics,
 266-274
 vs nesiritide (ROSE trial), 266-274, 270-273
 Dopamine receptor agonists, 165t
 Doppler echocardiography, 28, 28t
 DOSE trial, 259-260, 261, 274
 DPP-4 inhibitors, 165-166, 165t
 Dyspnea, 21, 22t, 56, 255, 256t
 nocturnal, 255
 vasodilators for, 262-264
 Edema, 255, 256t
 acute pulmonary, 39t, 43, 256t
 pulmonary edema, 22t, 43
 Ejection fraction. See also *Heart failure with preserved ejection fraction (HF-pEF)*; *Heart failure with reduced ejection fraction (HF-rEF)*.
 heart failure with mid-range ejection fraction (HF-mEF),
 xviii
 Ejection fraction (*continued*)
 heart failure with reduced ejection fraction (HF-rEF),
 xviii, 38-39
 treatment recommendations, xviii
 left ventricular (LVEF), 97, 110, 185
 in HF pathophysiology, 38
 treatment decisions and, xviii
 Elderly patients. See also *Psychosocial issues*.
 atrial fibrillation in, 127, 128
 cognitive impairment/dementia, 282-283
 frailty, 279-282
 heart failure in, 37t
 Electrocardiogram, 24-27, 26t
 12-lead, 24-27, 26t
 common findings in HF, 26t
 Electrolytes, 25t
 EMPA-REG OUTCOME, 174-175
 Empagliflozin, 165t
 Enalapril, 92-93, 189-190
 End-of-life planning, 288-291
 clinical indications for (ESC Guidelines), 288t
 management plan, discussing with patients and families,
 289-291, 290t
 palliative care, 288-289, 289
 suggestions for communication with patients, 289-291, 290t
 Endomyocardial biopsy, 31
 Endothelin receptor antagonists, 132t
 Entresto. See *Sacubitril/valsartan*.
 Environmental conditions, 78-79
 altitude, 78
 particulate pollution, 79
 temperature, 78-79
 Eplerenone, 102
 clinical trials, 103, 106-107
 EPHESUS, 103
 dosage, 104t-105t
 indications for, 103
 ESCAPE trial, 39, 55
 Escitalopram, xx
 EVEREST trials, 133, 260-262, 263, 274
 hospitalizations and future risk, 254

Exercise, 77, 287-288
cardiopulmonary exercise evaluation, 30t
clinical trials of, 287
depression, reduction in symptoms, 287-288
in HF-pEF, 77
in HF-rEF, 77
Exercise intolerance
in diagnosis, 21, 22t
onset of, 22-23
Exercise testing, 30, 30t

FAIR-HF study, 193-194, 195, 202
Familial heart disease, 22, 23t
Fibrillation, atrial. See *Atrial fibrillation*.
Filling pressures, 31, 39t, 50-51, 51, 57
in HF-pEF, 68t
Fluid restriction, 72
Fluid retention, 255, 256t. See also *Edema*.
FRAIL-HF study, 280-282
Frailty, 279-282
ADHF patients, 280-281
diagnosis of, 279-280
gait speed, 279-280
PRISMA-7 survey, 279-280
ESC guideline recommendations on, 280t
FRAIL-HF, 280-282
medication review, 280t
referral to specialist, 280t
Furosemide, 33, 115
vs torsemide, 122t-124t

Gender, hospitalization rates for HF and, 34, 35, 36t
GISSI-HF study, 76
Glimepiride, 165t
Glipizide, 165t
Glitazones, 165t
Glomerular filtration rate (GRF), 25t, 61, 104t-105t
Glyburide, 165t
Guidelines for heart failure management, xiv-xx
2013 ACCF/AHA Guideline, xiv
2016 ACC/AHA/HFSA, xiv-xv
2016 ESC Guidelines, xiv, xv-xx, 258-259

Guidelines for heart failure management (*continued*)
ACCF/AHA recommendation system, xvi-xviii[t], 258
consensus view in, xx
critical information on, xiv
ESC Guidelines, xiv, xv-xx, 258-259
on psychosocial issues, 280t, 281t
guideline-directed therapy, xii, xiv
for ivabradine, xix, xv
level (quality) of evidence, xvii[t]
limitations of, xii
for sacubitril/valsartan, xv
strength/class of recommendations, xvi[t]
updates to, xiv-xx

Heart failure. See also *Treatment of heart failure*.
acute heart failure, 253-277
classification of, 33-47
clinical syndromes and staging, 33-47
comorbidities, 36t-37t, 159-216, 160t, 162t-163t
diagnosis of, 21-32
hemodynamic patterns in, 39-40, 39t
interventions for, 217-252
pathophysiology of, 49-69
psychosocial issues, 279-294
resources for patients, 295-296
subcategories of, 22t. See also *specific subcategories*.
heart failure with mid-range ejection fraction (HF-mEF), xviii
heart failure with preserved ejection fraction (HF-pEF), 56-61
heart failure with reduced ejection fraction (HF-rEF), xviii, 38-39
timing of: acute vs chronic, 22t
symptoms of, 21, 22t, 256t-257t
Heart failure with mid-range ejection fraction (HF-mEF), xviii
Heart failure with preserved ejection fraction (HF-pEF), 44t
acute decompensated heart failure (ADHF) in, 253
comorbidities, 49, 66-67
chronic kidney disease (CKD), 190-193, 191, 192, 193
COPD, 60
definition of, 56-57

Heart failure with preserved ejection fraction (HF-pEF)
(continued)

- exercise recommendations, 77
- pathophysiology of, 56-61, 64
 - cardiac changes, 68t
 - comorbidities and, 49, 60, 66-67
 - hypertension, 88
 - musculoskeletal responses, 61, 68t
 - pulmonary responses, 57-60, 68t
 - renal responses, 60-61, 67
- phenotype, 43, 49
- physiology of (LV pressure/volume), 56-57, 62-63
- precapillary pulmonary hypertension and, 131-132
- treatment
 - ACEi-s, 8
 - ARBs, 8
 - β -blockers, 98-99
 - hydralazine/ISDN, 113
 - MRAs, 106-109
 - sacubitil/valsartan, 91

Heart failure with reduced ejection fraction (HF-rEF), 38-39, 44t

- acute decompensated heart failure (ADHF) in, 253
- comorbidities
 - anemia, 195
 - treatment algorithms, 201, 202
 - chronic kidney disease (CKD), 186, 187t, 189-190, 192
 - COPD, 173-179
- effectiveness of therapies, xviii, 49
- exercise recommendations, 77, 287-288
- pathophysiology in, 49-56
 - CAD and, 161
 - cardiovascular responses, 51-54, 52-53
 - hemodynamic congestion, 50-51
 - myocardial injury as initiator, 49, 50
 - negative feedback cycles, 49-56, 50, 83-84
 - pulmonary abnormalities, 54-55
 - RAAS activation, 50, 50, 51, 51
 - renal responses, 55-56, 55t, 56t
 - salt and water retention, 50, 50
- treatment
 - ACEi-s, xviii, 84-86

Heart failure with reduced ejection fraction (HF-rEF), treatment *(continued)*

- ARBs, 86-88
- β -blockers, xviii, 96
- digoxin, 117
- inotropes, 274
- ivabradine, 99
- MRAs, xviii, 102-106
- sacubitil/valsartan, 91

Heart rate

- CardioMEMS HF System for monitoring, 237
- in diagnosis, 24t, 26t
 - elevated, 50
 - inotropic agents and, 267t
 - ivabradine for heart rate reduction, xix, 101, 173-179, 184t, 185
 - resting, 26t
 - sympathetic nervous system and, 50
- Heart rhythm, 26t
- Heart transplant, 30, 31, 73, 237
- HeartMate II device, 242, 243

Hemodynamics

- in acute decompensated heart failure
 - deterioration, as marker for risk of adverse outcomes, 264-266
 - deterioration preceding clinical events, 253
 - hemodynamic congestion, 50-51
 - hemodynamic patterns in HF, 39-40, 39t
 - in advanced HF, 244t-245t
 - in cardiogenic shock, 39t, 244t
 - implantable hemodynamic monitoring, 231-237
 - vasodilator use and, 264-266, 265t
- Hemoglobin/hematocrit, 25t
- HF-ACTION Study, 287
 - patient phenotypes (“clusters”) in, 46-47
- HF-pEF. See *Heart failure with preserved ejection fraction*.
- HF-rEF. See *Heart failure with reduced ejection fraction*.
- History, in diagnosis, 22-23
 - cardiovascular history, 23, 23t
 - familial heart disease, 22, 23t

Hospitalization for heart failure
for acute decompensated heart failure (ADHF), 253-254,
254, 264-266, 265t
demographic data, 36t
discharges, number of, 34, 34
hospitalization rates, trends in, 34-35, 34, 36t
hospitalizations for depression, 283-287, 286
hypotension and, 264, 265t
mortality risk and, 253-254, 254, 264, 265t
before and after discharge, 254
number of HF hospitalizations and, 254, 254
Hydralazine, 109-113
action mechanism, 109-111
African Americans and, 111-113, 112
clinical trials of
 A-HeFT, 111-113, 112
 V-HeFT-I and V-HeFT-II, 111
in combination with ISDN, 111, 112-113
hemodynamic effects, 110-111
indications for, 111, 112-113
Hydrochlorothiazide, 113, 114t
Hyperkalemia, 102
ACE inhibitors and, 86t
ARBs and, 86t
CKD patients, 194t
MRAs and, 106, 109
Hyperlipidemia, 160t
Hypertension, 38, 60, 131-133. See also *Diuretics*.
alcohol intake and, 74, 75, 76
chronic pulmonary, 38, 60
heart failure and, 36t, 160t
precapillary pulmonary hypertension, 131-132
pulmonary, agents targeting, 131-133
Hypokalemia, diuretics and, 115
Hyponatremia, 257t
 diuretics and, 115
 fluid restriction for, 72
Hypotension
 ACE inhibitors and, 86t
 ARBs and, 86t
 hospitalization and, 264, 265t
 vasodilators and, 264, 265t

Imaging studies, in diagnosis of HF, 28-29, 29t
2D and Doppler echocardiography, 28, 28t
Implantable electrophysiologic devices.
See *Cardiovascular implantable electrophysiologic devices*.
Implantable hemodynamic monitoring, 231-237
CardioMEMS HF System, 236-237
 FDA approval of, 237
CHAMPION trial, 236-237
 changes in medications, 236-237, 240
 dosage changes in medications, 237, 241
 management algorithm, 237, 238-239
COMPASS-HF trial, 231-236
Individuals
 individualization of therapy, 255, 274
 prognosis, variation in, xiii
Infection
 ADHF precipitated by, 259t
 implanted electrophysiologic devices (CIEDs) and,
 152-153, 154-155
Inotropes, 266-274, 267t
 adverse events, 266, 268t-269t
 agents available, 267t
 indications for, 274
 acute decompensated HF-rEF, 274
 cardiogenic shock, 274
 hospitalized patients with severe systolic dysfunction, 274
 low-dose, 266-274, 270-273
 not recommended for HF-rEF patients with ADHF, 266
INTERMACS registry of HF stages, 242, 244t-245t, 248
Interventions for HF, 217-252
 aortic stenosis, 217-219, 218, 220-223
 implantable hemodynamic monitoring, 231-237
 left atrial appendage (LAA) closure, 228-231
 mechanical circulatory support, 237-246
 mitral regurgitation, surgical repair of, 219-228
 transcatheter aortic valve replacement (TAVR), 217-219
Iron/iron-binding capacity, 25t
Iron deficiency, 193
 algorithm for diagnosis, 201
 algorithm for treatment, 202
Iron supplementation, 194-195, 196-200

Ischemic heart disease, comorbid with HF, 160t
Ivabradine, 99-102
action mechanism, 99
 β -blockers and, xv
clinical trials, SHIFT, 99, 100-101
dosage, initiation and up-titration of, xv
efficacy, 99-102, 100-101
FDA approval, 99
guidelines for, xv
indications for, xix, 99-101
2016 ACC/AHA/HFSA guidelines, xv
EMA approved uses, xix
heart rate reduction in HF-rEF patients with COPD, xix,
101, 173-179, 184t, 185
vs β -blockers, 99

Kidney disease. See *Chronic kidney disease (CKD)*.
Kinins, 84

Laboratory tests, 23-24, 25
LCZ696, xviii, 89. See also *Sacubitril/valsartan*.
Left atrial appendage (LAA) closure, 228-231
AF patients with contraindication for chronic
anticoagulation, 228-229, 231
Kaplan-Meier curves, 229, 232-233
Kaplan-Meier curves for bleeding, 234-235
Watchman Device, 229-231, 230

Left ventricular assist devices (LVADs), 237-242. See also
Mechanical circulatory support.
patient eligibility criteria, 246, 251t

Left ventricular ejection fraction (LVEF), xviii, 97, 110, 185
Left ventricular (left heart) failure, 38-39, 38
impaired LV systolic function (HF-rEF), 38
Leukocytes/leukocytosis, 25t
Linagliptin, 165t
Loop diuretics, 115-116. See also *Diuretics*.
torsemide, 115-116, 118-121
torsemide vs furosemide, 116, 122t-124t, 125t

LVEF. See *Left ventricular ejection fraction*.

Magnetic resonance imaging (MRI)
Mechanical circulatory support, 237-246
ECS Guidelines, 246, 248-249
for patient eligibility, 246, 251t
HeartMate II device, 242, 243
INTERMACS registry of advanced HF stages, 242,
244t-245t, 248
left ventricular assist devices (LVADs), 237-242, 248-249
patient eligibility criteria, 246, 251t
REMATCH study, 237
ROADMAP study, 242-246, 243, 249
terms used in indications for, 250t
treatment algorithm and decision guide, 248-249
Meglitinides, 165t
Mental status assessment, 281t, 282-283, 284-285
Metformin, 165t, 166
Metoprolol, 194t
Metoprolol succinate (sustained-release), 96-97
Microalbuminuria, 186
Miglitol, 165t
Milrinone, 266, 267t, 268t-269t
adverse events, 266, 268t-269t
Mineralcorticoid receptor antagonists (MRAs), 102-109.
See also *specific agents*.
action mechanism, 102
adverse events, 106
hyperkalemia, 106, 109
renal dysfunction, 109
approved agents (eplerone, spironolactone), 102
clinical trials, 103-109
EPHESUS, 103
RALES, 103
TOPCAT, 107, 108
combined with chlorthalidone (diuretic), 115
discontinuation of, 103
dosing, 104t-105t
efficacy, 103-109
guidelines and recommendations, xviii, 102, 109
indications for
HF-pEF, 106-109
HF-rEF, xviii, 102-106
HF with CKD, 194t
monitoring, 103, 109, 110t

- Mitral regurgitation (MR), 133, 219-228
ADHF precipitated by, 259t
catheter-based approach, 225
EVEREST II trial, 225-226
exercise and, 54
guidelines on repair, 219-228
in HF-rEF, 54
incidence of, 219
MitraClip device, 133, 225-228, 225
complications, 228t
indications for, 226
meta-analysis of outcomes, 226, 227
safety data, 226-228
surgical repair of, 219-228
Mitral valve regurgitation, 35
MOOD-HF Study, 287
MRAs. See *Mineralcorticoid receptor antagonists*.
Musculoskeletal responses to HF, 56, 60t, 61, 68t
nitrate supplements and, 76-77
Myocardial fibrosis, torsemide effects, 120-121
Myocardial infarction, prior, heart failure and, 36t, 38
Myocardial injury, 35
in HF-rEF pathophysiology, 49, 50
Myocardial ischemia, stress testing for, 30t
Myocardial remodeling. See *Remodeling*.
Myocardium
contractility of, 52-53
“stretch-squeeze” relationship, 52, 54
- Nateglinide, 165t
Natriuretic peptides, 88
A-type (ANP), 51, 88
augmentation of, 88
B-type (BNP), 51, 88, 169, 179, 256t
RAAS activation and, 51, 51
using to evaluate HF with COPD, 169, 179
Negative feedback cycles, 49-56, 50, 83-84
in HF and CKD, 186, 189
Neprilysin enzyme, xv, 88. See also *ARNI (angiotensin receptor neprilysin inhibitor)*.
- Neprilysin (NEP) inhibitor. See also *Sacubitril/valsartan*.
angiotensin receptor neprilysin inhibitor (ARNI), xv, 88
LCZ696 and, xviii, 89
Nesiritide, 264, 265t, 266-274
low-dose (“renal-dose”), to enhance effects of diuretics, 266-274
vs dopamine (ROSE trial), 266-274, 270-273
Neuro-hormonal antagonists (ACEi-s, MRAs, β-blockers), xviii
New York Heart Association (NYHA) classification system, 40-43
ACC/AHA stages, relationship to, 42
maximal oxygen uptake, 40, 40
- Nitrates, 109-113
action mechanism, 109-111
dietary supplementation, 76-77
hemodynamic effects, 110-111
hydralazine, 109-113
Nitric oxide, 76-77
Nitroglycerine, 264
Nitroprusside, 264
hemodynamic effects of, 111
Nonpharmacologic management of chronic heart failure, 71-82. See also *Diet*.
dietary interventions, 71-77
environmental conditions, 78-79
physical activity, 77
NSAIDS, ADHF precipitated by, 259t
NYHA classification. See *New York Heart Association (NYHA) classification system*.
- Obesity, 67, 73
comorbid with HF, 162t
ONTARGET, 87
OPTIME-CHF, 266, 268t-269t, 274
Orthopnea, 255, 258
Oxygen, maximal consumption of, in HF-pEF, 57
Oxygen saturation, low, 256t

- Pacing
 cardiac resynchronization therapy, 143, 146, 148-149
 in AF patients, 147-151
 dyssynchrony induced by RV pacing, 147, 150, 150t
 right-ventricular, 150t
- Palliative care, 288-289, 289
- PARADIGM-HF, 89, 92-93
- PARAMOUNT trial, 60-61, 91-96, 96
- Particulate pollution, 79
- Pathophysiology of heart failure, 49-69
 in HF-pEF, 56-61, 64
 cardiac changes, 68t
 comorbidities, 49, 66-67
 musculoskeletal responses, 61, 68t
 physiology: LV volume/pressure, 62-63
 pulmonary responses, 57-60, 68t
 renal responses, 60-61, 67
 in HF-rEF, 49-56, 50
 cardiac performance and end diastolic volume, 52-53
 cardiovascular responses, 51-54, 52-53
 hemodynamic congestion, 50-51
 musculoskeletal responses, 56, 60t
 negative feedback cycles, 49-56, 50, 83-84
 pulmonary abnormalities, 54-55
 RAAS activation, 50, 50, 51, 51
 renal responses, 55-56, 55t, 56t
 salt and water retention, 50, 50
 self-perpetuating process, 51
- Patient phenotypes (“clusters”) in HF, 43, 46-47
- Patient questions, information needed to address, 33
- Patients and families
 discussions about prognoses, 289, 290t
 end-of-life planning, 288-291
 management plan, discussing with, 289-291, 290t
- Peripheral perfusion, assessment of, 23, 24t, 258
- Peripheral vascular disease, heart failure and, 36t
- Pharmacologic management of heart failure, 83-140, 258-274. See also *specific drugs and drug types*.
 acute decompensated heart failure (ADHF), 255, 258-274
 agents, 84t
 ACEI-s, 83-86
 agents targeting pulmonary hypertension, 131-133, 132t
- Pharmacologic management of heart failure, agents (*continued*)
 anti-platelet agents, 125-127
 anti-thrombotic agents, 125-131
 anticoagulants, 127-131
 ARBs, 86-88
 β-blockers, 96-99
 digoxin, 116-117, 126
 diuretics, 113-116, 259-260
 dobutamine, 267t
 dopamine, 266-274, 267t
 hydralazine/nitrates, 109-113
 inotropes, 266-274, 267t
 ivabradine, 99-102
 milrinone, 266, 267t
 morphine, 258-259
 MRAs (mineralcorticoid receptor antagonists), 102-109
 sacubitril/valsartan, 88-96
 vasodilators, 109-113, 262-266, 274
 vasopressin antagonists, 260-262, 263
 of chronic heart failure, 83-140
 classes of agents, 84t
- Phenotypes of HF patients, 43, 46-47
- Phosphodiesterase inhibitors
 indications for, 132t
 milrinone, 266, 267t
- Physical activity, 77. See also *Exercise*.
- Physical examination, 23, 24t
- Pioglitazone, 165t
- Prazosin (Minipress), 111
- Prognosis
 depression and, 283, 286
 discussions with patients and families about, 289, 290t
 individual variation in, xiii
- Prostanoids, 132t
- Psychosocial issues, 37t, 279-294
 cognitive impairment, 281t, 282-283, 284-285
 depression, xix-xx, 283-288
 end-of-life planning, 288-291, 288t
 ESC guidelines on, 280t, 281t
 frailty, 279-282, 280t
- Pulmonary abnormalities, in HF-rEF, 54-55

Pulmonary congestion, 256t
Pulmonary edema, 22t
 acute, patient phenotype, 43
 hemodynamic patterns in, 39t
Pulmonary examination, 23, 24t
Pulmonary function testing, 29-30, 30t
Pulmonary hypertension
 agents targeting, 131-133, 132t
 referral to a HF center, 132
Pulmonary responses
 in HF-pEF, 57-60, 68t
 in HF-rEF, 54-55

QRS widening, 143, 146
QT-prolongation, anticancer drugs and, 203

RAAS. See *Renin-angiotensin-aldosterone system*.
Race
 hospitalization rates for HF and, 35, 36t
 hydralazine, African Americans and, 111-113, 112
RALES trial, 103
Ramipril, clinical trials of, 87
RED-HF trial, 193
Regurgitation. See *Mitral valve regurgitation*.
REMATCH study, 237
Remodeling, 35
 electrical, 54
 in HF-rEF, 50, 54
 negative feedback cycle, 50
 structural, 54
Renal dysfunction, 67
 cardio-renal syndrome, 55, 55t
 heart failure and, 37t
 in HF-pEF, 60-61, 67
 in HF-rEF, 55-56, 55t, 56t
 mechanisms of, 55-56, 56t
 pathophysiology of, 55-56, 56t, 58-59
 renal congestion, 55-56, 58
 renal inflammation, 58-59
Renal function
 glomerular filtration rate, 25t, 104t-105t
 monitoring with MRAs, 109
 testing for, 25t

Renin, increased levels of, 50
Renin-angiotensin-aldosterone system (RAAS), 50, 50
 activation of, 50, 50
 cascade of effects, 85
 sodium (salt) limitation and, 71
 activation of natriuretic peptides, 51, 51
 aldosterone in, 102
 renin-angiotensin cascade, 85
 torsemide effects on, 118-119
Repaglinide, 165t
Resources for patients, 295-296
Right heart (Swan-Ganz) catheterization, 31, 39
Right ventricular failure, 35-38, 38
Right-ventricular pacing, cardiac effects of, 150t
Riociguat, 132t
Rivaroxaban, 130-131
ROADMAP study, 242-246, 243, 249
ROSE trial, 266-274, 270-273, 274
Rosiglitazone, 165t

Sacubitril
 action mechanism, 89
 as LCZ696, xviii, 89
Sacubitril/valsartan (Entresto), 88-96
 action mechanism, xv, 89
 adverse reactions, xv
 angioedema, xv, 89
 minimizing risk for, xviii
 hypotension, xv
 cautions and contraindications, xv, 89
 combining with ACEi or ARB, xviii
 clinical trials, 89-96
 PARADIGM-HF, 89, 92-93
 PARAMOUNT, 91-96, 96
 dosing
 initiation of therapy, 91, 94-95
 switching from ACEi-s or ARBs to, 91, 94-95
 efficacy, 89-96
 FDA approval, 89
 guidelines for, xv, 91
 indications for, xv
 2016 ACC/AHA/HFSA guidelines, xv, 91

Sacubitril/valsartan (Entresto), indications for (*continued*)
HF-pEF, 91
HF-rEF, 91
to replace ACEi in HF-rEF patients, xviii, 91, 94
initiation of, withholding ACEi prior to, xviii
LCZ696, xviii, 89
molecular structure, 90
troponin T-sensitivity, 92, 96
SADHART-CHF, 287
Salt limitation (in diet), 71-72, 72t
Salt retention, 21, 22t, 50, 50
Saxagliptin, 165t
Selixigap, 132t
Sertraline, 287
SGLT2 inhibitors, 165t, 167-169, 168
adverse effects of, 169, 176-177
CV outcomes trials, 178t
SHIFT trial, 99, 100-101
Shock
cardiogenic, 22t, 39t
hemodynamic patterns in, 39t
Sildenafil, 132t
Silent cerebral ischemia (SCI), 282, 283
Sitagliptin, 165t
Sleep-related breathing disorders, 23t, 67
Sodium limitation, 71-72
conversion factors, 72t
Sodium retention, ADHF precipitated by, 259t
SOLVD, 84
renal function in, 189-190
Spironolactone, 102
clinical trials, 103, 106-107
RALES, 103
TOPCAT, 107, 108
dosage, 104t-105t
indications for, 103, 106, 109
ST-T waves, 26t
Staging of heart failure, 33-47. See also *Classification of heart failure*.
ACC/AHA stages, 40-43, 41t
INTERMACS registry of stages for advanced HF, 242, 244t-245t
NYHA classification system, 40-43, 40, 41t, 42

Statistically significant P-values, xii
STICHES, 159-161, 164
“Stretch-squeeze” relationship, 52, 54
Stroke, heart failure and, 36t, 127
Substance abuse, 23t, 43
ADHF precipitated by, 259t
Sudden cardiac death, 141-143
incidence of, 142
Sulfonylureas, 165t
Surgical treatments. See also *Interventions for HF*.
aortic stenosis, 217-219, 218, 220-223
cardiac transplantation, 30, 31, 73, 237
cardiovascular implantable electrophysiologic devices (CIEDs), 141-158
implantable hemodynamic monitoring, 231-237
left atrial appendage (LAA) closure, 228-231
mechanical circulatory support, 237-246
mitral regurgitation, surgical repair of, 219-228
Sympathetic nervous system, 50
Symptoms of heart failure, 21, 22t, 254-255, 256t-257t
T waves, 26t
Tachycardia, 26t, 50
low cardiac output and, 257t
milrinone, 268t
sudden cardiac death and, 141, 142
Tamponade, MitraClip and, 228t
Telmisartan (Micardis), 87
Temperature extremes (environmental), 78-79
Thiazide diuretics, 113-115
Thiazolidinediones
adverse effects of, 161
in diabetes, 161
Thromboses: anti-thrombotic agents, 125-131
Thyroid function tests, 25t
Tolvaptan, 260-262, 263
approved uses, 262
TOPCAT, 107, 108
Torsemide, 115-116
effects on myocardial fibrosis, 120-121
effects on RAAS, 118-119
potential benefits of, 125t
vs furosemide, 122t-124t, 125t

Transcatheter aortic valve replacement (TAVR), 217-219
Treatment of heart failure. See also *specific subjects*.
in acute decompensated heart failure (ADHF), 255, 258-274
AV nodal ablation, 147-151, 151
cardiac resynchronization therapy, 143, 146, 148-149
cardiovascular implantable electrophysiologic devices, 141-158
classification and staging systems and, 33-47
patient phenotypes, 43, 46-47
treatment context provided, 43, 44t
goals of, xviii, 255
guidelines for, xiv-xx, 258-259. See also *Guidelines for heart failure management*.
2016 ESC Guidelines, xiv, xv-xx, 258-259
on psychosocial issues, 280t, 281t
ACC/AHA guidelines, xvi-xviii[t], 258
guideline-directed therapy, xii, xiv
limitations of, xii
neuro-hormonal antagonists (ACEi-s, MRAs, β -blockers), xviii
statistical P-values, xii
strength of recommendations and level of evidence, xvi-xviii[t]
updated guidelines, xiv-xx
interventions for HF, 217-252
aortic stenosis, 217-219
implantable hemodynamic monitoring, 231-237
mechanical circulatory support, 237-246
nonpharmacologic management, 71-82
dietary interventions, 71-77
environmental conditions, 78-79
physical activity, 77
palliative care, 288-289, 289
pharmacologic management, 83-140, 84t
ACEi-s, 84-86
agents targeting pulmonary hypertension, 131-133, 132t
anti-platelet agents, 125-127
anti-thrombotic agents, 125-131
anticoagulants, 127-131
ARBs, 86-88
 β -blockers, 96-99
of chronic heart failure, 83-140

Treatment of heart failure, pharmacologic management (*continued*)
digoxin, 116-117, 126
diuretics, 113-116, 259-260
dobutamine, 267t
dopamine, 266-274, 267t
hydralazine/nitrates, 109-113
inotropes, 266-274, 267t
ivabradine, 99-102
milrinone, 266, 267t
mineralcorticoid receptor antagonists (MRAs), 102-109
morphine, 258-259
sacubitri/valsartan, 88-96
vasodilators, 109-113, 262-266, 274
vasopressin antagonists, 260-262, 263
psychosocial issues, 279-294
cognitive impairment, 282-283
depression, 283-288
end-of-life planning, 288-291
frailty, 279-282
QRS widening, 143, 146
Tricyclic antidepressants, xx
Troponin T, 92, 96
Two-dimensional echocardiography, 28, 28t
Type 2 diabetes, 161-169, 165t, 170-172
U waves, 26t
Ultrafiltration, 262, 274
Valsartan. See also *Sacubitri/valsartan*.
molecular structure, 90
Vascular resistance, systemic, 50
Vasodilators, 109-113, 262-266, 274. See also *specific drugs*.
ACC/AHA guidelines, 264
blood pressure, concerns about, 264, 265t
choice of agent, 264
hypotension and, 264, 265t
nesiritide, 264, 265t
nitroglycerine, 264
nitroprusside, 264
parenteral, 262-264
use in acute decompensated heart failure, 262-266

Vasopressin antagonists, 260-262, 263
tolvaptan, 260-262, 263

Ventricle
left
left ventricular assist devices (LVADs), 237-242
left ventricular ejection fraction (LVEF), xviii, 97, 110, 185
left ventricular failure, 38-39, 38
physiology in HF-pEF: LV pressure/volume, 62-63
right
right ventricular failure, 35-38, 38
right-ventricular pacing, 150t

Ventricular dysfunction, in heart failure, 52-53, 54

Ventricular remodeling. See *Remodeling*.

Volume overload, 22t, 50
causes of, 50, 50

Volume status assessment, 23, 24t

Warfarin, 127, 128. See also *Anticoagulants*.
use with clopidogrel, 130
Watchman Device as alternative to, 229

Watchman Device, 229-231, 230

Water restriction. See *Fluid restriction*.

Water retention, 50, 50

Weight gain, 23t

Weight loss, recommendation for, 73-74

White blood cell count, 25t